

KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2313**Specification****KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	P42574
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 32 kDa , observed, 35 kDa KDa
Gene Name	CASP3
Aliases	CASP3; Caspase 3; Apopain; CPP32; CPP32B; Caspase 3, Apoptosis-Related Cysteine Peptidase; Caspase 3, Apoptosis-Related Cysteine Protease; SREBP Cleavage Activity 1; Cysteine Protease CPP32; Protein Yama; EC 3.4.22.56; Caspase-3; CASP-3; CPP-32; SCA-1; Yama; PARP Cleavage Protease; Procaspase3; EC 3.4.22
Immunogen	A synthesized peptide derived from human pro Caspase 3

KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody - Additional Information

Gene ID	836
Other Names	
Caspase-3, CASP-3, 3.4.22.56, Apopain, Cysteine protease CPP32, CPP-32, Protein Yama, SREBP cleavage activity 1, SCA-1, Caspase-3 subunit p17, Caspase-3 subunit p12, CASP3, CPP32 {ECO:0000303 PubMed:7983002}	

KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody - Protein Information**Name** CASP3**Synonyms** CPP32 {ECO:0000303|PubMed:7983002}**Function**

Thiol protease that acts as a major effector caspase involved in the execution phase of apoptosis (PubMed:18723680, PubMed:20566630, PubMed:23650375, PubMed:35338844, PubMed:35446120, PubMed:7596430). Following cleavage and activation by initiator caspases (CASP8, CASP9 and/or CASP10), mediates

execution of apoptosis by catalyzing cleavage of many proteins (PubMed:18723680, PubMed:20566630, PubMed:23650375, PubMed:7596430). At the onset of apoptosis, it proteolytically cleaves poly(ADP-ribose) polymerase PARP1 at a '216-Asp-I-Gly-217' bond (PubMed:10497198, PubMed:16374543, PubMed:7596430, PubMed:7774019). Cleaves and activates sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain (By similarity). Cleaves and activates caspase-6, -7 and -9 (CASP6, CASP7 and CASP9, respectively) (PubMed:7596430). Cleaves and inactivates interleukin-18 (IL18) (PubMed:37993714, PubMed:9334240). Involved in the cleavage of huntingtin (PubMed:8696339). Triggers cell adhesion in sympathetic neurons through RET cleavage (PubMed:21357690). Cleaves DSG2 in response to apoptosis resulting in a loss of full length DSG2 at desmosome cell junctions and subsequent loss of cell-cell adhesion (PubMed:17559062). Also cleaves JUP in response to apoptosis (PubMed:17559062). Cleaves and inhibits serine/threonine-protein kinase AKT1 in response to oxidative stress (PubMed:23152800). Acts as an inhibitor of type I interferon production during virus-induced apoptosis by mediating cleavage of antiviral proteins CGAS, IRF3 and MAVS, thereby preventing cytokine overproduction (PubMed:30878284). Also involved in pyroptosis by mediating cleavage and activation of gasdermin-E (GSDME) (PubMed:35338844, PubMed:35446120). Cleaves XRCC4 and phospholipid scramblase proteins XKR4, XKR8 and XKR9, leading to promote phosphatidylserine exposure on apoptotic cell surface (PubMed:23845944, PubMed:33725486). Cleaves BIRC6 following inhibition of BIRC6-caspase binding by DIABLO/SMAC (PubMed:36758104, PubMed:36758106).

Cellular Location

Cytoplasm.

Tissue Location

Highly expressed in lung, spleen, heart, liver and kidney. Moderate levels in brain and skeletal muscle, and low in testis. Also found in many cell lines, highest expression in cells of the immune system.


KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-pro Caspase 3 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-pro Caspase 3 antibody (Cat#AGI2313). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-pro Caspase 3 antibody (Cat#AGI2313, 1:10,000) and HRP-conjugated goat anti rabbit secondary antibody respectively.



Western blotting analysis using anti-pro Caspase 3 antibody (Cat#AGI2313). pro Caspase 3 expression in wild type (WT) and pro Caspase 3 shRNA knockdown (KD) HeLa cells with 30 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-pro Caspase 3 antibody (Cat#AGI2313, 1:10,000) and HRP-conjugated goat anti rabbit secondary antibody respectively.



Flow cytometric analysis of pro Caspase 3 expression in HT-1080 cells using pro Caspase 3 antibody (Cat#AGI2313,1:2,000). Green, isotype control; red, pro Caspase 3.



Immunocytochemical staining of C2C12 cells with pro Caspase 3 antibody (Cat#AGI2313, 1:1,000). Nuclei were stained blue with DAPI; Pro Caspase 3 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.