

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2322

Specification

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody - Product Information

Application	WB, FC, ICC
Primary Accession	Q96KQ7
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 132 kDa; Observed, 160-170 kDa
Gene Name	KDa
Aliases	EHMT2 Euchromatic Histone Lysine Methyltransferase 2; KMT1C; G9A; C6orf30; BAT8; Euchromatic Histone-Lysine N-Methyltransferase 2; Histone-Lysine N-Methyltransferase EHMT2; Histone H3-K9 Methyltransferase 3; HLA-B Associated Transcript 8; Lysine N-Methyltransferase 1C; H3-K9-HMTase 3; Em:AF134726.3; NG36/G9a; NG36; Histone-Lysine N-Methyltransferase, H3 Lysine-9 Specific 3; Chromosome 6 Open Reading Frame 30; Ankyrin Repeat-Containing Protein; G9A Histone Methyltransferase; HLA-B-Associated Transcript 8; EC 2.1.1.367; Protein G9a; EC 2.1.1.-; GAT8
Immunogen	A synthesized peptide derived from human EHMT2

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody - Additional Information

Gene ID	10919
Other Names	
Histone-lysine N-methyltransferase EHMT2, 2.1.1.-, 2.1.1.367, Euchromatic histone-lysine N-methyltransferase 2, HLA-B-associated transcript 8, Histone H3-K9 methyltransferase 3, H3-K9-HMTase 3, Lysine N-methyltransferase 1C, Protein G9a, EHMT2, BAT8, C6orf30, G9A, KMT1C, NG36	

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody - Protein Information

Name EHMT2

Synonyms BAT8, C6orf30, G9A, KMT1C, NG36

Function

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones. Also mediates monomethylation of 'Lys-56' of histone H3 (H3K56me1) in G1 phase, leading to promote interaction between histone H3 and PCNA and regulating DNA replication. Also weakly methylates 'Lys-27' of histone H3 (H3K27me). Also required for DNA methylation, the histone methyltransferase activity is not required for DNA methylation, suggesting that these 2 activities function independently. Probably targeted to histone H3 by different DNA-binding proteins like E2F6, MGA, MAX and/or DP1. May also methylate histone H1. In addition to the histone methyltransferase activity, also methylates non-histone proteins: mediates dimethylation of 'Lys-373' of p53/TP53. Also methylates CDYL, WIZ, ACIN1, DNMT1, HDAC1, ERCC6, KLF12 and itself.

Cellular Location

Nucleus. Chromosome. Note=Associates with euchromatic regions (PubMed:11316813). Does not associate with heterochromatin (PubMed:11316813).

Tissue Location


Expressed in all tissues examined, with high levels in fetal liver, thymus, lymph node, spleen and peripheral blood leukocytes and lower level in bone marrow

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-EHMT2 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-EHMT2 antibody (Cat#68352). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with


anti-EHMT2 antibody (Cat#68352, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-EHMT2 antibody (Cat#68352). EHMT2 expression in wild type (WT) and EHMT2 shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-EHMT2 antibody (Cat#68352, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).



Flow cytometric analysis of EHMT2 expression in HepG2 cells using EHMT2 antibody (Cat#68352, 1:2,000). Green, isotype control; red, EHMT2.



Immunocytochemical staining of HepG2 cells with EHMT2 antibody (Cat#68352, 1:1,000). Nuclei were stained blue with DAPI; EHMT2 was stained magenta with Alexa Fluor® 647. Scale bar: 20 µm.