

KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2327

Specification

KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody - Product Information

| | |
|-------------------|---|
| Application | WB, FC, ICC |
| Primary Accession | 075955 |
| Reactivity | Rat, Human, Mouse |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Calculated MW | Predicted, 47 kDa ; Observed, 48 kDa KDa |
| Gene Name | FLOT1 |
| Aliases | Flotillin 1; Flotillin-1; Integral Membrane Component Of Caveolae |
| Immunogen | A synthesized peptide derived from human Flotillin 1 |

KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody - Additional Information

| | |
|-------------|--------------------|
| Gene ID | 10211 |
| Other Names | Flotillin-1, FLOT1 |

KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody - Protein Information

Name FLOT1

Function

May act as a scaffolding protein within caveolar membranes, functionally participating in formation of caveolae or caveolae-like vesicles.

Cellular Location

Cell membrane; Peripheral membrane protein. Endosome Membrane, caveola {ECO:0000250|UniProtKB:O08917}; Peripheral membrane protein {ECO:0000250|UniProtKB:O08917}. Melanosome. Membrane raft. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Membrane-associated protein of caveola (By similarity) {ECO:0000250|UniProtKB:O08917, ECO:0000269|PubMed:17081065}


KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-Flotillin 1 Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-Flotillin 1 antibody (Cat#AGI2327). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Flotillin 1 antibody (Cat#AGI2327, 1:10,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.




Western blotting analysis using anti-Flotillin 1 antibody (Cat#AGI2327). Flotillin 1 expression in wild type (WT) and Flotillin 1 shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-Flotillin 1 antibody (Cat#AGI2327, 1:10,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Copyright ©2024 Genus Biotechnologies LLC

Flow cytometric analysis of Flotillin 1 expression in HepG2 cells using anti-Flotillin 1 antibody (Cat#AGI2327, 1:2,000). Green, isotype control; red, Flotillin 1.



Copyright ©2024 Genus Biotechnologies LLC

Immunocytochemical staining of HepG2 cells with Flotillin 1 antibody (Cat#AGI2327, 1:1,000). Nuclei were stained blue with DAPI; Flotillin 1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μ m.