

KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI2347**Specification****KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	P61158
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 47 kDa; Observed, 47 kDa
Gene Name	KDa ACTR3
Aliases	ACTR3; Actin Related Protein 3; ARP3; ARP3 Actin Related Protein 3 Homolog; Actin-Related Protein 3; Actin-Like Protein 3; ARP3 (Actin-Related Protein 3, Yeast) Homolog; ARP3 Actin-Related Protein 3 Homolog (Yeast)
Immunogen	A synthesized peptide derived from human ACTR3

KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody - Additional Information

Gene ID	10096
Other Names	
Actin-related protein 3, Actin-like protein 3, ACTR3, ARP3	

KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody - Protein Information**Name** ACTR3**Synonyms** ARP3**Function**

ATP-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF) (PubMed:9000076). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility (PubMed:9000076). Seems to contact the pointed end of the daughter actin filament (PubMed:9000076). In podocytes, required for the formation of lamellipodia downstream of AVIL and PLCE1 regulation (PubMed:29058690). In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:17220302, PubMed:29925947).

The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed:29925947). Plays a role in ciliogenesis (PubMed:20393563).

Cellular Location


Cytoplasm, cytoskeleton. Cell projection. Nucleus. Note=In pre- apoptotic cells, colocalizes with MEKV in large specks (pyroptosomes) (PubMed:19109554)

KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-Arp3 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-Arp3 antibody (Cat#69173). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Arp3 antibody (Cat#69173, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:50,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-Arp3 antibody (Cat#69173). Arp3 expression in wild type (WT) and Arp3 shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-Arp3 antibody (Cat#69173, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:50,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).



Flow cytometric analysis of Arp3 expression in C2C12 cells using Arp3 antibody (Cat#69173, 1:2,000). Green, isotype control; red, Arp3.



Immunocytochemical staining of C2C12 cells with Arp3 antibody (Cat#69173, 1:1,000). Nuclei were stained blue with DAPI; Arp3 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.