

**Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone SPM165 ]**  
**Catalog # AH10726****Specification**

---

**Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Product Information**

Application	WB, IHC-P, IF, FC
Primary Accession	<a href="#">Q07817</a>
Other Accession	<a href="#">598</a> , <a href="#">516966</a>
Reactivity	Human, Mouse, Rat, Pig
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2a
Calculated MW	27kDa kDa

**Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Additional Information****Gene ID** 598**Other Names**

Bcl-2-like protein 1, Bcl2-L-1, Apoptosis regulator Bcl-X, BCL2L1, BCL2L, BCLX

**Application Note**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_FC">FC~~1:10~50</span>

**Format**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &amp; 0.05% azide. Also available WITHOUT BSA &amp; azide at 1.0mg/ml.

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Protein Information****Name** BCL2L1**Synonyms** BCL2L, BCLX**Function**

Potent inhibitor of cell death. Inhibits activation of caspases. Appears to regulate cell death by blocking the voltage- dependent anion channel (VDAC) by binding to it and preventing the release of the caspase activator, CYC1, from the mitochondrial membrane. Also acts as a regulator of G2

checkpoint and progression to cytokinesis during mitosis. Isoform Bcl-X(S) promotes apoptosis.

#### Cellular Location

[Isoform Bcl-X(L)]: Mitochondrion inner membrane. Mitochondrion outer membrane Mitochondrion matrix. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Cytoplasm, cytosol. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus membrane; Single-pass membrane protein; Cytoplasmic side. Note=After neuronal stimulation, translocates from cytosol to synaptic vesicle and mitochondrion membrane in a calmodulin-dependent manner (By similarity). Localizes to the centrosome when phosphorylated at Ser-49

#### Tissue Location


Bcl-X(S) is expressed at high levels in cells that undergo a high rate of turnover, such as developing lymphocytes. In contrast, Bcl-X(L) is found in tissues containing long-lived postmitotic cells, such as adult brain

### Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Hodgkin's Lymphoma stained with Bcl-x Monoclonal Antibody (SPM165).

### Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 27kDa, identified as the Bcl-X protein. This MAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241 amino acid protein which suppresses cell death. And Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain which functions as a dominant inhibitor of Bcl-2. This MAb reacts with both Bcl-XS and Bcl-XL proteins.

**Bcl-X (Apoptosis Marker) Antibody - With BSA and Azide - References**

Hsu YT, et. al. Journal of Biological Chemistry, 1997, 272(21):13829-34. | Hsu YT, et. al. Proceedings of the National Academy of Sciences of the United States of America, 1997, 94(8):3668-72. | Wolter KG, et. al. Journal of Cell Biology, 1997, 139(5):1281-92