

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone CDH16/1071 ]**  
**Catalog # AH10983**

---

**Specification**

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide -  
Product Information**

Application	IHC, IF, FC
Primary Accession	<a href="#">O75309</a>
Other Accession	<a href="#">1014</a> , <a href="#">513660</a>
Reactivity	Human, Mouse, Rat, Rabbit, Dog
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	130kDa KDa

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide -  
Additional Information**

**Gene ID** 1014

**Other Names**

Cadherin-16, Kidney-specific cadherin, Ksp-cadherin, CDH16

**Application Note**

IHC~~1:100~500  
IF~~1:50~200  
FC~~1:10~50

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide -  
Protein Information**

**Name** CDH16

**Function**

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein


**Tissue Location**

Kidney specific.


**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Images**

Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with KSP-Cadherin Monoclonal Antibody (CDH16/1071)



Formalin-fixed, paraffin-embedded Rat Kidney stained with KSP-Cadherin Monoclonal Antibody (CDH16/1071)

**Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - Background**

This MAb recognizes a protein of 130kDa, identified as Ksp-cadherin. Cadherins form a superfamily

of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins have been implicated in embryogenesis, tissue morphogenesis, tissue structure maintenance, cell polarization, neoplastic invasiveness and metastasis, and membrane transport. It is suggested that Ksp-cadherin is a marker for terminal differentiation of the basolateral membranes of renal tubular epithelial cells. Within the kidney, Ksp-Cadherin is found exclusively in the basolateral membrane of renal tubular epithelial cells and collecting duct cells, and not in glomeruli, renal interstitial cells, or blood vessels. Ksp-Cadherin has been suggested to distinguish Chromophobe Renal-Cell Carcinoma from Oncocytoma.

#### **Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody - With BSA and Azide - References**

Shen, S.S., et al. 2005. Kidney-specific cadherin, a specific marker for the distal portion of the nephron and related renal neoplasms. *Mod Pathol.* 18: 933-40. | Mazal, P.R., et al. 2005. Expression of kidney-specific cadherin distinguishes chromophobe renal cell carcinoma from renal oncocytoma. *Hum Pathol.* 36: 22-28. |