

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide**Mouse Monoclonal Antibody [Clone MBS-12]****Catalog # AH11148****Specification**

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Product Information

Application	IHC, IF, FC
Primary Accession	P02771
Other Accession	174 , 518808
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	70kDa KDa

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Additional Information**Gene ID** 174**Other Names**

Alpha-fetoprotein, Alpha-1-fetoprotein, Alpha-fetoglobulin, AFP, HPAFP

Application Note

IHC~~1:100~500
IF~~1:50~200
FC~~1:10~50

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Protein Information**Name** AFP**Synonyms** HPAFP**Function**

Binds copper, nickel, and fatty acids as well as, and bilirubin less well than, serum albumin. Only a small percentage (less than 2%) of the human AFP shows estrogen-binding properties.

Cellular Location

Secreted.


Tissue Location

Plasma. Synthesized by the fetal liver and yolk sac

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Images

Formalin-fixed, paraffin-embedded human Fetal Liver stained with AFP Monoclonal Antibody (MBS-12).

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - Background

It recognizes an oncofetal glycoprotein with a single chain of 70kDa, which is identified as alpha fetoprotein (AFP). This MAb is highly specific to AFP and shows no cross-reaction with other oncofetal antigens or serum albumin. The yolk sac and the liver produce AFP during fetal life. AFP expression in adults is often associated with hepatoma or teratoma. However, hereditary persistence of alpha-fetoprotein may also be found in individuals with no obvious pathology. The protein is thought to be the fetal counterpart of serum albumin, and the AFP and albumin genes are present in tandem in the same transcriptional orientation on chromosome 4. AFP is found in monomeric as well as dimeric and trimeric forms, and binds copper, nickel, fatty acids and bilirubin. The level of AFP in amniotic fluid is used to measure renal loss of protein to screen for spinal bifida and anencephaly.Å

AFP (Alpha Fetoprotein) (Hepatocellular/Germ Cell Tumor Marker) Antibody - With BSA and Azide - References

Stefanova, I., Horejs♦, V., Kristofov♦, H., Angelisov♦, P., Zizkovský, V. and Hilgert, I. 1988. Monoclonal antibodies against human α -fetoprotein. Exploitation of an unusual calcium-dependent interaction with the antigen for analytical and preparative purposes. J. Immunol. Methods 111: 67-73. | Lafuste, P., et al. 2002. α -fetoprotein gene expression in early and full-term human trophoblast. Placenta 23: 600-612