

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide
Rabbit Polyclonal Antibody [Clone]
Catalog # AH11497**Specification****IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Product Information**

Application	WB, IF, FC
Primary Accession	075874
Other Accession	3417 , 593422
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit / Immunoglobulin
Calculated MW	45-47kDa KDa

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Additional Information**Gene ID** 3417**Other Names**

Isocitrate dehydrogenase [NADP] cytoplasmic, IDH, 1.1.1.42, Cytosolic NADP-isocitrate dehydrogenase, IDP, NADP(+)-specific ICDH, Oxalosuccinate decarboxylase, IDH1, PICD

Application Note

WB~~1:1000
IF~~1:50~200
FC~~1:10~50

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Protein Information**Name** IDH1**Synonyms** PICD**Function**

Catalyzes the NADP(+)-dependent oxidative decarboxylation of isocitrate (D-threo-isocitrate) to 2-ketoglutarate (2-oxoglutarate), which is required by other enzymes such as the phytanoyl-CoA dioxygenase (PubMed: [10521434](http://www.uniprot.org/citations/10521434)), PubMed: [19935646](http://www.uniprot.org/citations/19935646)). Plays a critical role in the generation of NADPH, an important cofactor in many biosynthesis pathways (PubMed: [10521434](http://www.uniprot.org/citations/10521434)). May act as a

corneal epithelial crystallin and may be involved in maintaining corneal epithelial transparency (By similarity).

Cellular Location


Cytoplasm, cytosol. Peroxisome

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Images



Western Blot Analysis of A431 Cell Lysate using IDH1 Polyclonal Antibody (Rabbit)

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - Background

It recognizes a 45kDa protein, which is identified as isocitrate dehydrogenase (IDH1). It belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 catalyzes the third step of the citric acid cycle, which involves the oxidative decarboxylation of isocitrate, forming α -ketoglutarate and CO_2 in a two-step reaction. The first step involves the oxidation of isocitrate to the intermediate oxalosuccinate, while the second step involves the production of α -ketoglutarate. During this process, either NADH or NADPH is produced along with CO_2 . Recently, an inactivating mutation of IDH1 has been implicated in glioblastoma. IDH1 appears to function as a tumor suppressor that, when mutationally inactivated, contributes to tumorigenesis in part through induction of the HIF-1 pathway.

IDH1 (Isocitrate Dehydrogenase) Antibody - With BSA and Azide - References

Geisbrecht, B.V. and Gould, S.J. 1999. The human PICD gene encodes a cytoplasmic and peroxisomal NADP⁺-dependent isocitrate dehydrogenase. J. Biol. Chem. 274: 30527-30533