

**Anti-NSE gamma Antibody**  
**Mouse Monoclonal Antibody**  
**Catalog # AH13196**

### Specification

#### Anti-NSE gamma Antibody - Product Information

Application	WB, IHC-P, IF, FC
Primary Accession	<a href="#">P09104</a>
Other Accession	<a href="#">511915</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b
Calculated MW	47269

#### Anti-NSE gamma Antibody - Additional Information

##### Gene ID 2026

##### Other Names

2-phospho-D-glycerate hydrolyase; ENO2; ENOG; Enolase 2 gamma neuronal; Enolase2; Gamma-enolase; Neural enolase; Neuron specific gamma enolase; Neuron-specific enolase; NSE

##### Application Note

<span class ="dilution\_WB">WB~~1:1000</span><br /><span class ="dilution\_IHC-P">IHC-P~~N/A</span><br /><span class ="dilution\_IF">IF~~1:50~200</span><br /><span class ="dilution\_FC">FC~~1:10~50</span>

##### Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

##### Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

##### Precautions

Anti-NSE gamma Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### Anti-NSE gamma Antibody - Protein Information

##### Name ENO2 ([HGNC:3353](#))

##### Function

Enolase that catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate in glycolysis and the reverse reaction in gluconeogenesis (By similarity). Has neurotrophic and neuroprotective properties on a broad spectrum of central nervous system (CNS) neurons. Binds, in a calcium-dependent manner, to cultured neocortical neurons and promotes cell survival (By

similarity).

#### Cellular Location

Cytoplasm. Cell membrane. Note=Can translocate to the plasma membrane in either the homodimeric (alpha/alpha) or heterodimeric (alpha/gamma) form

#### Tissue Location


The alpha/alpha homodimer is expressed in embryo and in most adult tissues. The alpha/beta heterodimer and the beta/beta homodimer are found in striated muscle, and the alpha/gamma heterodimer and the gamma/gamma homodimer in neurons

### Anti-NSE gamma Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-NSE gamma Antibody - Images



Formalin-fixed, paraffin-embedded Human Pheochromocytoma stained with NSE gamma Monoclonal Antibody (SPM347).

### Anti-NSE gamma Antibody - Background

Recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha-isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphoenolpyruvic acid in the glycolytic pathway.

NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It is usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.