

Anti-MART-1 / Melan-A / MLANA Antibody
Recombinant Rabbit Monoclonal Antibody
Catalog # AH13229

Specification

Anti-MART-1 / Melan-A / MLANA Antibody - Product Information

Application	WB, IHC-P, IF, FC
Primary Accession	Q16655
Other Accession	154069
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal
Isotype	Rabbit / IgG, kappa
Calculated MW	13157

Anti-MART-1 / Melan-A / MLANA Antibody - Additional Information

Gene ID 2315

Other Names

Antigen LB39-AA, Antigen SK29-AA, Melanoma antigen recognized by T-cells 1, MLAN-A, MLANA

Application Note

WB~~1:1000
IHC-P~~N/A
IF~~1:50~200
FC~~1:10~50

Format

200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Anti-MART-1 / Melan-A / MLANA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-MART-1 / Melan-A / MLANA Antibody - Protein Information

Name MLANA

Synonyms MART1

Function

Involved in melanosome biogenesis by ensuring the stability of GPR143. Plays a vital role in the expression, stability, trafficking, and processing of melanocyte protein PMEL, which is critical to the formation of stage II melanosomes.

Cellular Location

Endoplasmic reticulum membrane; Single-pass type III membrane protein. Golgi apparatus. Golgi apparatus, trans-Golgi network membrane. Melanosome. Note=Also found in small vesicles and tubules dispersed over the entire cytoplasm. A small fraction of the protein is inserted into the membrane in an inverted orientation Inversion of membrane topology results in the relocalization of the protein from a predominant Golgi/post-Golgi area to the endoplasmic reticulum. Melanoma cells expressing the protein with an inverted membrane topology are more effectively recognized by specific cytolytic T-lymphocytes than those expressing the protein in its native membrane orientation


Tissue Location

Expression is restricted to melanoma and melanocyte cell lines and retina


Anti-MART-1 / Melan-A / MLANA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MART-1 / Melan-A / MLANA Antibody - Images

Formalin-fixed, paraffin-embedded Human Skin stained with MART-1 Recombinant Rabbit Monoclonal Antibody (MLANA/1761R)



Formalin-fixed, paraffin-embedded Human Melanoma stained with MART-1 Recombinant Rabbit Monoclonal Antibody (MLANA/1761R)

Anti-MART-1 / Melan-A / MLANA Antibody - Background

This antibody recognizes a protein doublet of 20-22kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Seven other melanoma associated antigens recognized by autologous cytotoxic T cells include MAGE-1, MAGE-3, tyrosinase, gp100, gp75, BAGE-1, and GAGE-1. Subcellular fractionation shows that MART-1 is present in melanosomes and endoplasmic reticulum. This MAb labels melanomas and other tumors showing melanocytic differentiation. It is also a useful positive-marker for angiomyolipomas. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin.