

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody
Mouse Monoclonal Antibody
Catalog # AH13432**Specification**

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Product Information

Application	WB, IHC-P, IF, FC
Primary Accession	P01009
Other Accession	525557
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1
Calculated MW	46737

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Additional Information**Gene ID** 5265**Other Names**

A1AT; AAT; Alpha 1 antiproteinase Alpha 1-antitrypsin; Alpha-1 protease inhibitor; Alpha-1-antiproteinase; alpha1 proteinase inhibitor; Alpha1AT; Dom1; Serine (or cysteine) proteinase inhibitor clade A member 1; Serine protease inhibitor 1-1; Serine protease inhibitor A1a; Serpin A1a; Serpin peptidase inhibitor clade A member 1; Serpina1; Short peptide from AAT; SPAAT; Spi1-1

Application Note

WB~~1:1000
IHC-P~~N/A
IF~~1:50~200
FC~~1:10~50

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Protein Information**Name** SERPINA1 ([HGNC:8941](#))**Synonyms** AAT, PI

Function

Inhibitor of serine proteases. Its primary target is elastase, but it also has a moderate affinity for plasmin and thrombin. Irreversibly inhibits trypsin, chymotrypsin and plasminogen activator. The aberrant form inhibits insulin-induced NO synthesis in platelets, decreases coagulation time and has proteolytic activity against insulin and plasmin.

Cellular Location

Secreted. Endoplasmic reticulum. Note=The S and Z allele are not secreted effectively and accumulate intracellularly in the endoplasmic reticulum


Tissue Location

Ubiquitous. Expressed in leukocytes and plasma.

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Images

Formalin-fixed, paraffin-embedded human Tonsil stained with Alpha-1-Antitrypsin Monoclonal Antibody (AAT/1379)

Anti-Alpha-1-Antitrypsin (SERPINA1) Antibody - Background

The immunohistochemical staining of AAT is useful in identification of benign and malignant hepatic tumors and yolk sac carcinomas. Positive staining for AAT is also used in detection of benign and malignant lesions of histiocytic nature. This antibody is may also useful tool in the screening of patients with cryptogenic cirrhosis or other forms of liver disease with fibrosis of uncertain origin.