

ASH2L antibody - middle region

Rabbit Polyclonal Antibody
Catalog # Al10117

Specification

ASH2L antibody - middle region - Product Information

Application WB, IHC Primary Accession Q9UBL3

Other Accession O9UBL3, NP_004665, NM_004674

Reactivity Human, Mouse, Rat, Rabbit, Dog, Guinea

Pig, Horse, Bovine

Predicted Human, Mouse, Rat, Chicken, Dog, Guinea

Pig, Horse, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 69 kDa KDa

ASH2L antibody - middle region - Additional Information

Gene ID 9070

Alias Symbol ASH2, ASH2L1, ASH2L2, Bre2

Other Names

Set1/Ash2 histone methyltransferase complex subunit ASH2, ASH2-like protein, ASH2L, ASH2L1

Target/Specificity

ASH2L is a component of the Set1/Ash2 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3, but not if the neighboring 'Lys-9' residue is already methylated. The protein may function as a transcriptional regulator and play a role in hematopoiesis.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ASH2L antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

ASH2L antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ASH2L antibody - middle region - Protein Information

Name ASH2L

Synonyms ASH2L1





Function

Transcriptional regulator (PubMed:12670868). Component or associated component of some histone methyltransferase complexes which regulates transcription through recruitment of those complexes to gene promoters (PubMed:19131338). Component of the Set1/Ash2 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3, but not if the neighboring 'Lys-9' residue is already methylated (PubMed:19556245). As part of the MLL1/MLL complex it is involved in methylation and dimethylation at 'Lys-4' of histone H3 (PubMed:19556245). May play a role in hematopoiesis (PubMed:12670868). In association with RBBP5 and WDR5, stimulates the histone methyltransferase activities of KMT2A, KMT2B, KMT2C, KMT2D, SETD1A and SETD1B (PubMed:21220120, PubMed:22266653).

Cellular Location

Nucleus.

Tissue Location


Ubiquitously expressed. Predominantly expressed in adult heart and testis and fetal lung and liver, with barely detectable expression in adult lung, liver, kidney, prostate, and peripheral leukocytes.

ASH2L antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ASH2L antibody - middle region - Images




ASH2L antibody - middle region (Al10117) in Human Placenta cells using Immunohistochemistry Immunohistochemistry with Human Placenta lysate tissue at an antibody concentration of 5.0µg/ml using anti-ASH2L antibody (Al10117)





ASH2L antibody - middle region (AI10117) in Human Testis cells using Immunohistochemistry Immunohistochemistry with Human Testis lysate tissue at an antibody concentration of 5.0µg/ml using anti-ASH2L antibody (AI10117)



ASH2L antibody - middle region (Al10117) in Human HeLa cells using Western Blot

WB Suggested Anti-ASH2L Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: Hela cell lysate

ASH2L antibody - middle region - Background

This is a rabbit polyclonal antibody against ASH2L. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).