

Twist1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11347**Specification**

Twist1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9EPJ1
Other Accession	NM_053530 , NP_445982
Reactivity	Human, Mouse, Rat, Zebrafish, Horse, Bovine
Predicted Host	Rat, Zebrafish, Pig, Chicken, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 22kDa KDa

Twist1 antibody - N-terminal region - Additional Information

Alias Symbol	Twist
--------------	-------

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Twist1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Twist1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Twist1 antibody - N-terminal region - Protein Information**Twist1 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Twist1 antibody - N-terminal region - Images



WB Suggested Anti-Twist1 Antibody Titration: 1.0 µg/ml
Positive Control: Rat Liver

Twist1 antibody - N-terminal region - Citations

- [Inhibition of ATM reverses EMT and decreases metastatic potential of cisplatin-resistant lung cancer cells through JAK/STAT3/PD-L1 pathway.](#)
- [Effects of secreted frizzled-related protein 1 on proliferation, migration, invasion, and apoptosis of colorectal cancer cells.](#)