

C6orf134 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12129**Specification**

C6orf134 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5SQI0
Other Accession	NM_024909 , NP_079185
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36kDa kDa

C6orf134 antibody - N-terminal region - Additional Information**Gene ID** 79969**Alias Symbol** DKFZp547J097, FLJ13158, Nbla00487, TAT, MEC17, C6orf134**Other Names**

Alpha-tubulin N-acetyltransferase 1 {ECO:0000255|HAMAP-Rule:MF_03130}, Alpha-TAT {ECO:0000255|HAMAP-Rule:MF_03130}, Alpha-TAT1 {ECO:0000255|HAMAP-Rule:MF_03130}, TAT {ECO:0000255|HAMAP-Rule:MF_03130}, 2.3.1.108 {ECO:0000255|HAMAP-Rule:MF_03130}, Acetyltransferase mec-17 homolog {ECO:0000255|HAMAP-Rule:MF_03130}, ATAT1 {ECO:0000255|HAMAP-Rule:MF_03130}

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C6orf134 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C6orf134 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

C6orf134 antibody - N-terminal region - Protein Information**Name** ATAT1 {ECO:0000255|HAMAP-Rule:MF_03130}**Function**

Specifically acetylates 'Lys-40' in alpha-tubulin on the luminal side of microtubules. Promotes microtubule destabilization and accelerates microtubule dynamics; this activity may be

independent of acetylation activity. Acetylates alpha-tubulin with a slow enzymatic rate, due to a catalytic site that is not optimized for acetyl transfer. Enters the microtubule through each end and diffuses quickly throughout the lumen of microtubules. Acetylates only long/old microtubules because of its slow acetylation rate since it does not have time to act on dynamically unstable microtubules before the enzyme is released. Required for normal sperm flagellar function. Promotes directional cell locomotion and chemotaxis, through AP2A2-dependent acetylation of alpha-tubulin at clathrin-coated pits that are concentrated at the leading edge of migrating cells. May facilitate primary cilium assembly.

Cellular Location


Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03130, ECO:0000269|PubMed:24097348}. Membrane, clathrin-coated pit {ECO:0000255|HAMAP-Rule:MF_03130, ECO:0000269|PubMed:24097348}. Cell junction, focal adhesion {ECO:0000255|HAMAP-Rule:MF_03130, ECO:0000269|PubMed:24097348}. Cell projection, axon {ECO:0000255|HAMAP-Rule:MF_03130}. Cytoplasm, cytoskeleton {ECO:0000255|HAMAP-Rule:MF_03130}. Cytoplasm, cytoskeleton, spindle {ECO:0000255|HAMAP-Rule:MF_03130}

C6orf134 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C6orf134 antibody - N-terminal region - Images



WB Suggested Anti-C6orf134 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:1562500

Positive Control: ACHN cell lysate