

RUNDC2A antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12830**Specification**

RUNDC2A antibody - N-terminal region - Product Information

Application	WB, IHC
Primary Accession	O9HA26
Other Accession	NM_032167 , NP_115543
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Dog
Predicted Host	Human, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 42kDa KDa

RUNDC2A antibody - N-terminal region - Additional Information

Alias Symbol **FLJ12363**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-RUNDC2A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


RUNDC2A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RUNDC2A antibody - N-terminal region - Protein Information**RUNDC2A antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RUNDC2A antibody - N-terminal region - Images



Human Intestine



WB Suggested Anti-RUNDC2A Antibody Titration: 5.0µg/ml
Positive Control: HepG2 cell lysate

RUNDC2A antibody - N-terminal region - References

Bonaldo, M.F., (1996) Genome Res. 6(9), 791-806 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.