

CTAG1B Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14559**Specification**

CTAG1B Antibody - middle region - Product Information

Application	WB
Primary Accession	P78358
Other Accession	NM_001327 , NP_001318
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18kDa KDa

CTAG1B Antibody - middle region - Additional Information**Gene ID** 1485;246100**Alias Symbol** **CT6.1, CTAG, CTAG1, CTAG1A, ESO1, LAGE-2, LAGE2B, NY-ESO-1****Other Names**

Cancer/testis antigen 1, Autoimmunogenic cancer/testis antigen NY-ESO-1, Cancer/testis antigen 6.1, CT6.1, L antigen family member 2, LAGE-2, CTAG1A, CTAG, CTAG1, ESO1, LAGE2, LAGE2A

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CTAG1B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CTAG1B Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

CTAG1B Antibody - middle region - Protein Information**Name** CTAG1A**Synonyms** CTAG, CTAG1, ESO1, LAGE2, LAGE2A**Cellular Location**

Cytoplasm

Tissue Location

Expressed in testis and ovary and in a wide variety of cancers. Detected in uterine myometrium. Expressed from 18 weeks until birth in human fetal testis. In the adult testis, is strongly expressed


in spermatogonia and in primary spermatocytes, but not in post-meiotic cells or in testicular somatic cells (at protein level)

CTAG1B Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CTAG1B Antibody - middle region - Images



WB Suggested Anti-CTAG1B Antibody Titration: 1.0 µg/ml
Positive Control: Fetal Stomach

CTAG1B Antibody - middle region - References

Chen Y.-T., et al. Proc. Natl. Acad. Sci. U.S.A. 94:1914-1918(1997).
Lethe B.G., et al. Int. J. Cancer 76:903-908(1998).
Wang R.-F., et al. J. Immunol. 161:3598-3606(1998).
De Smet C., et al. Mol. Cell. Biol. 19:7327-7335(1999).
Aradhya S., et al. Hum. Mol. Genet. 10:2557-2567(2001).