

**FGFR1OP2 Antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15269****Specification**

---

**FGFR1OP2 Antibody - C-terminal region - Product Information**

|                   |                                                                |
|-------------------|----------------------------------------------------------------|
| Application       | WB                                                             |
| Primary Accession | <a href="#">Q9NVK5</a>                                         |
| Other Accession   | <a href="#">NM_015633</a> , <a href="#">NP_056448</a>          |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Predicted         | Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog |
| Host              | Rabbit                                                         |
| Clonality         | Polyclonal                                                     |
| Calculated MW     | 29kDa KDa                                                      |

**FGFR1OP2 Antibody - C-terminal region - Additional Information****Gene ID** 26127**Alias Symbol** DKFZp564O1863, DKFZp586C1423, FLJ37569, HSPC123-like, WIT3.0**Other Names**

FGFR1 oncogene partner 2, FGFR1OP2

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-FGFR1OP2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

FGFR1OP2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**FGFR1OP2 Antibody - C-terminal region - Protein Information****Name** FGFR1OP2**Function**

May be involved in wound healing pathway.

**Cellular Location**

Cytoplasm.

**Tissue Location**


Expressed in bone marrow, spleen and thymus.

### **FGFR1OP2 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **FGFR1OP2 Antibody - C-terminal region - Images**



WB Suggested Anti-FGFR1OP2 Antibody Titration: 1.0 µg/ml  
Positive Control: MCF7 Whole Cell

### **FGFR1OP2 Antibody - C-terminal region - References**

Zhang Q.-H., et al. Genome Res. 10:1546-1560(2000).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Lin L., et al. Submitted (DEC-2003) to the EMBL/GenBank/DDBJ databases.  
Bechtel S., et al. BMC Genomics 8:399-399(2007).  
Grand E.K., et al. Genes Chromosomes Cancer 40:78-83(2004).