

PTGER3 / EP3 Antibody (Cytoplasmic Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10228**Specification**

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Product Information

Application	IHC-P
Primary Accession	P43115
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	43kDa KDa
Dilution	IHC-P~~N/A

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Additional Information**Gene ID** 5733**Other Names**

Prostaglandin E2 receptor EP3 subtype, PGE receptor EP3 subtype, PGE2 receptor EP3 subtype, PGE2-R, Prostanoid EP3 receptor, PTGER3

Target/Specificity

Human PTGER3 / EP3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

PTGER3 / EP3 Antibody (Cytoplasmic Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Protein Information**Name** PTGER3**Function**

Receptor for prostaglandin E2 (PGE2) (PubMed:7883006, PubMed:7981210, PubMed:8117308, PubMed:8135729, PubMed:8307176). The activity of this receptor can couple to both the inhibition of adenylate cyclase mediated by G(i) proteins, and to an elevation of intracellular calcium (PubMed:7883006, PubMed:7981210, PubMed:8117308, PubMed:8135729, PubMed:8307176).

href="http://www.uniprot.org/citations/8117308" target="_blank">8117308, PubMed:8135729). Required for normal development of fever in response to pyrinogens, including IL1B, prostaglandin E2 and bacterial lipopolysaccharide (LPS). Required for normal potentiation of platelet aggregation by prostaglandin E2, and thus plays a role in the regulation of blood coagulation. Required for increased HCO₃(-) secretion in the duodenum in response to mucosal acidification, and thereby contributes to the protection of the mucosa against acid- induced ulceration. Not required for normal kidney function, normal urine volume and osmolality (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Detected in kidney (PubMed:8117308, PubMed:8135729). Expressed in small intestine, heart, pancreas, gastric fundic mucosa, mammary artery and pulmonary vessels


Volume

50 µl


PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Images


Anti-PTGER3 / EP3 antibody IHC of human Breast, Carcinoma.




Human, Breast: Formalin-Fixed Paraffin-Embedded (FFPE)



Anti-PTGER3 / EP3 antibody IHC of human brain, hippocampus.



Anti-PTGER3 / EP3 antibody IHC of human brain, hippocampus.



Anti-PTGER3 / EP3 antibody IHC of human Pancreas, Carcinoma.

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - Background

Receptor for prostaglandin E2 (PGE2); the EP3 receptor may be involved in inhibition of gastric acid secretion, modulation of neurotransmitter release in central and peripheral neurons, inhibition of sodium and water reabsorption in kidney tubulus and contraction in uterine smooth muscle. The activity of this receptor can couple to both the inhibition of adenylate cyclase mediated by G-I proteins, and to an elevation of intracellular calcium. The various isoforms have identical ligand binding properties but can interact with different second messenger systems (By similarity).

PTGER3 / EP3 Antibody (Cytoplasmic Domain) - References

- Adam M.,et al.FEBS Lett. 338:170-174(1994).
- Schmid A.,et al.Eur. J. Biochem. 228:23-30(1995).
- Yang J.,et al.Biochem. Biophys. Res. Commun. 198:999-1006(1994).
- Kunapuli S.P.,et al.Biochem. J. 298:263-267(1994).
- An S.,et al.Biochemistry 33:14496-14502(1994).