

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10726****Specification**

---

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC-P                  |
| Primary Accession | <a href="#">P25101</a> |
| Reactivity        | Human, Monkey          |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 49kDa KDa              |
| Dilution          | IHC-P ~ ~ N/A          |

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Additional Information****Gene ID** 1909**Other Names**

Endothelin-1 receptor, Endothelin A receptor, ET-A, ETA-R, hET-AR, EDNRA, ETA, ETRA

**Target/Specificity**

Human Endothelin A Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except EDNRB (72%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Protein Information****Name** EDNRA ([HGNC:3179](#))**Synonyms** ETA, ETRA**Function**

Receptor for endothelin-1. Mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system. The rank order of binding affinities for ET-A is: ET1 &gt; ET2 &gt;&gt; ET3.

**Cellular Location**

Cell membrane; Multi-pass membrane protein.

**Tissue Location**

Isoform 1, isoform 3 and isoform 4 are expressed in a variety of tissues, with highest levels in the

aorta and cerebellum, followed by lung, atrium and cerebral cortex, lower levels in the placenta, kidney, adrenal gland, duodenum, colon, ventricle and liver but no expression in umbilical vein endothelial cells. Within the placenta, isoform 1, isoform 2, isoform 3 and isoform 4 are expressed in the villi and stem villi vessels.


**Volume**

50 µl

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Images**

Anti-Endothelin A Receptor antibody ALS10726 IHC of human heart.

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - Background**

Receptor for endothelin-1. Mediates its action by association with G proteins that activate a phosphatidylinositol- calcium second messenger system. The rank order of binding affinities for ET-A is: ET1 > ET2 >> ET3.

**EDNRA / Endothelin A Receptor Antibody (Extracellular Domain) - References**

Adachi M.,et al.Biochem. Biophys. Res. Commun. 180:1265-1272(1991).  
Cyr C.,et al.Biochem. Biophys. Res. Commun. 181:184-190(1991).  
Hosoda K.,et al.FEBS Lett. 287:23-26(1991).  
Arai H.,et al.Jpn. Circ. J. 56:1303-1307(1992).  
Elshourbagy N.A.,et al.J. Biol. Chem. 268:3873-3879(1993).