

PPARG / PPAR Gamma Antibody (Internal)
Rabbit Polyclonal Antibody
Catalog # ALS11521**Specification**

PPARG / PPAR Gamma Antibody (Internal) - Product Information

Application	WB, IHC-P, E
Primary Accession	P37231
Reactivity	Human, Mouse, Rat, Rabbit, Hamster, Squirrel, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

PPARG / PPAR Gamma Antibody (Internal) - Additional Information**Gene ID** 5468**Other Names**

Peroxisome proliferator-activated receptor gamma, PPAR-gamma, Nuclear receptor subfamily 1 group C member 3, PPARG, NR1C3

Target/Specificity

Amino acids 255 -268 of human PPAR gamma isoform 1.

Reconstitution & Storage

+4°C or -20°C, Avoid repeated freezing and thawing.

Precautions

PPARG / PPAR Gamma Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

PPARG / PPAR Gamma Antibody (Internal) - Protein Information**Name** PPARG**Synonyms** NR1C3**Function**

Nuclear receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the nuclear receptor binds to DNA specific PPAR response elements (PPRE) and modulates the transcription of its target genes, such as acyl-CoA oxidase. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis. ARF6 acts as a key regulator of the tissue-specific adipocyte P2 (aP2) enhancer. Acts as a critical regulator of gut homeostasis by suppressing

NF-kappa-B-mediated pro-inflammatory responses. Plays a role in the regulation of cardiovascular circadian rhythms by regulating the transcription of BMAL1 in the blood vessels (By similarity).

Cellular Location

Nucleus. Cytoplasm. Note=Redistributed from the nucleus to the cytosol through a MAP2K1/MEK1-dependent manner. NOCT enhances its nuclear translocation

Tissue Location

Highest expression in adipose tissue. Lower in skeletal muscle, spleen, heart and liver. Also detectable in placenta, lung and ovary.


Volume

50 μ l


PPARG / PPAR Gamma Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPARG / PPAR Gamma Antibody (Internal) - Images

Anti-PPARG antibody IHC of human heart.



Anti-PPARG antibody IHC of human adrenal.



Anti-PPARG antibody IHC of human skin.

PPARG / PPAR Gamma Antibody (Internal) - Background

Nuclear receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Once activated by a ligand, the nuclear receptor binds to DNA specific PPAR response elements (PPRE) and modulates the transcription of its target genes, such as acyl-CoA oxidase. It therefore controls the peroxisomal beta-oxidation pathway of fatty acids. Key regulator of adipocyte differentiation and glucose homeostasis. ARF6 acts as a key regulator of the tissue-specific adipocyte P2 (aP2) enhancer. Acts as a critical regulator of gut homeostasis by suppressing NF-kappa-B-mediated proinflammatory responses. Plays a role in the regulation of cardiovascular circadian rhythms by regulating the transcription of ARNTL/BMAL1 in the blood vessels (By similarity).

PPARG / PPAR Gamma Antibody (Internal) - References

- Mukherjee R.,et al.J. Biol. Chem. 272:8071-8076(1997).
Elbrecht A.,et al.Biochem. Biophys. Res. Commun. 224:431-437(1996).
Yanase T.,et al.Biochem. Biophys. Res. Commun. 233:320-324(1997).
Greene M.E.,et al.Gene Expr. 4:281-299(1995).
Greene M.E.,et al.Submitted (DEC-2001) to the EMBL/GenBank/DDBJ databases.