

**TLR9 Antibody (Internal)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS11524****Specification**

---

**TLR9 Antibody (Internal) - Product Information**

Application	WB, IHC-P, IF, ICC
Primary Accession	<a href="#">Q9NR96</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	116kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A

**TLR9 Antibody (Internal) - Additional Information****Gene ID** 54106**Other Names**

Toll-like receptor 9, CD289, TLR9

**Target/Specificity**

15 amino acids from an internal region of human TLR9

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

**Precautions**

TLR9 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

**TLR9 Antibody (Internal) - Protein Information****Name** TLR9**Function**

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides (PubMed:<a href="http://www.uniprot.org/citations/14716310" target="\_blank">14716310</a>). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/11564765" target="\_blank">11564765</a>, PubMed:<a href="http://www.uniprot.org/citations/17932028" target="\_blank">17932028</a>). Controls lymphocyte response to Helicobacter infection (By

similarity). Upon CpG stimulation, induces B-cell proliferation, activation, survival and antibody production (PubMed:<a href="http://www.uniprot.org/citations/23857366" target="\_blank">23857366</a>).

#### Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9EQU3}. Early endosome membrane. Lysosome {ECO:0000250|UniProtKB:Q9EQU3} Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:Q9EQU3}. Golgi apparatus membrane. Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist. Exit from the ER requires UNC93B1. Endolysosomal localization is required for proteolytic cleavage and subsequent activation Intracellular localization of the active receptor may prevent from responding to self nucleic acid. {ECO:0000250|UniProtKB:Q9EQU3, ECO:0000269|PubMed:14716310, ECO:0000269|PubMed:38169466}

#### Tissue Location

Highly expressed in spleen, lymph node, tonsil and peripheral blood leukocytes, especially in plasmacytoid pre-dendritic cells. Levels are much lower in monocytes and CD11c+ immature dendritic cells. Also detected in lung and liver

#### Volume


50 µl

#### TLR9 Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### TLR9 Antibody (Internal) - Images




Immunocytochemistry of TLR9 in Jurkat cells with TLR9 antibody at 2 ug/ml.



Immunofluorescence of TLR9 in Mouse Spleen cells with TLR9 antibody at 10 ug/ml.



Western blot of TLR9 in Jurkat cell lysate with TLR9 antibody at (A) 0.5, (B) 1 and (C) 2 ug/ml.



Anti-TLR9 antibody IHC of human lung.

#### **TLR9 Antibody (Internal) - Background**

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine

secretion and the inflammatory response. Controls lymphocyte response to Helicobacter infection.

#### **TLR9 Antibody (Internal) - References**

Du X.,et al.Eur. Cytokine Netw. 11:362-371(2000).  
Chuang T.-H.,et al.Eur. Cytokine Netw. 11:372-378(2000).  
Hemmi H.,et al.Nature 408:740-745(2000).  
Liu Z.,et al.Submitted (SEP-2007) to the EMBL/GenBank/DDBJ databases.  
Nakajima T.,et al.Immunogenetics 60:727-735(2008).