

Apg7 / ATG7 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS12165**Specification**

Apg7 / ATG7 Antibody (C-Terminus) - Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	O95352
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A

Apg7 / ATG7 Antibody (C-Terminus) - Additional Information**Gene ID** 10533**Other Names**

Ubiquitin-like modifier-activating enzyme ATG7, ATG12-activating enzyme E1 ATG7, Autophagy-related protein 7, APG7-like, hAGP7, Ubiquitin-activating enzyme E1-like protein, ATG7, APG7L

Target/Specificity

17 amino acid peptide from near the carboxy terminus of human APG7.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

Apg7 / ATG7 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Apg7 / ATG7 Antibody (C-Terminus) - Protein Information**Name** ATG7 ([HGNC:16935](#))**Synonyms** APG7L**Function**

E1-like activating enzyme involved in the 2 ubiquitin-like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Facilitates LC3-I lipidation with

phosphatidylethanolamine to form LC3-II which is found on autophagosomal membranes (PubMed:34161705). Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Also plays a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation. Plays a role in regulating the liver clock and glucose metabolism by mediating the autophagic degradation of CRY1 (clock repressor) in a time-dependent manner (By similarity).

Cellular Location

Cytoplasm. Preautophagosomal structure. Note=Also localizes to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme

Tissue Location


Widely expressed, especially in kidney, liver, lymph nodes and bone marrow.

Apg7 / ATG7 Antibody (C-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Apg7 / ATG7 Antibody (C-Terminus) - Images



Anti-ATG7 antibody IHC of human liver.



Immunocytochemistry of APG7 in MCF7 cells with APG7 antibody at 10 ug/ml.



Immunofluorescence of APG7 in MCF7 cells with APG7 antibody at 20 ug/ml.

Apg7 / ATG7 Antibody (C-Terminus) - Background

E1-like activating enzyme involved in the 2 ubiquitin- like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Plays also a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation.

Apg7 / ATG7 Antibody (C-Terminus) - References

Yuan W.,et al.Mol. Biol. Cell 10:1353-1366(1999).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Muzny D.M.,et al.Nature 440:1194-1198(2006).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Tanida I.,et al.J. Biol. Chem. 276:1701-1706(2001).