

**GPI Antibody (N-Terminus)**  
**Goat Polyclonal Antibody**  
**Catalog # ALS13391****Specification**

---

**GPI Antibody (N-Terminus) - Product Information**

|                   |                                    |
|-------------------|------------------------------------|
| Application       | WB, IHC-P, E                       |
| Primary Accession | <a href="#">P06744</a>             |
| Reactivity        | Human, Monkey                      |
| Host              | Goat                               |
| Clonality         | Polyclonal                         |
| Calculated MW     | 63kDa KDa                          |
| Dilution          | WB~~1:1000<br>IHC-P~~N/A<br>E~~N/A |

**GPI Antibody (N-Terminus) - Additional Information****Gene ID** 2821**Other Names**

Glucose-6-phosphate isomerase, GPI, 5.3.1.9, Autocrine motility factor, AMF, Neuroleukin, NLK, Phosphoglucose isomerase, PGI, Phosphohexose isomerase, PHI, Sperm antigen 36, SA-36, GPI

**Target/Specificity**

Human GPI.

**Reconstitution & Storage**

Store at -20°C. Minimize freezing and thawing.

**Precautions**

GPI Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**GPI Antibody (N-Terminus) - Protein Information****Name** GPI {ECO:0000303|PubMed:2387591, ECO:0000312|HGNC:HGNC:4458}**Function**

In the cytoplasm, catalyzes the conversion of glucose-6-phosphate to fructose-6-phosphate, the second step in glycolysis, and the reverse reaction during gluconeogenesis (PubMed:<a href="http://www.uniprot.org/citations/28803808" target="\_blank">28803808</a>). Besides its role as a glycolytic enzyme, also acts as a secreted cytokine: acts as an angiogenic factor (AMF) that stimulates endothelial cell motility (PubMed:<a href="http://www.uniprot.org/citations/11437381" target="\_blank">11437381</a>). Acts as a neurotrophic factor, neuroleukin, for spinal and sensory neurons (PubMed:<a href="http://www.uniprot.org/citations/11004567" target="\_blank">11004567</a>, PubMed:<a href="http://www.uniprot.org/citations/3352745" target="\_blank">3352745</a>). It is secreted

by lectin-stimulated T-cells and induces immunoglobulin secretion (PubMed:<a href="http://www.uniprot.org/citations/11004567" target="\_blank">11004567</a>, PubMed:<a href="http://www.uniprot.org/citations/3352745" target="\_blank">3352745</a>).

#### Cellular Location

Cytoplasm. Secreted

#### GPI Antibody (N-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### GPI Antibody (N-Terminus) - Images



GPI antibody (0.3 ug/ml) staining of Human Heart lysate (35 ug protein in RIPA buffer).



Anti-GPI antibody IHC of human thyroid.

#### GPI Antibody (N-Terminus) - Background

Besides its role as a glycolytic enzyme, mammalian GPI can function as a tumor-secreted cytokine and an angiogenic factor (AMF) that stimulates endothelial cell motility. GPI is also a neurotrophic factor (Neuroleukin) for spinal and sensory neurons.

#### **GPI Antibody (N-Terminus) - References**

Gurney M.E., et al. Submitted (MAR-1987) to the EMBL/GenBank/DDBJ databases.  
Yakirevich E., et al. Biol. Reprod. 62:1016-1023(2000).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Grimwood J., et al. Nature 428:529-535(2004).  
Walker J.I.H., et al. Genomics 7:638-643(1990).