

MER / MERTK Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS13420**Specification**

MER / MERTK Antibody (C-Terminus) - Product Information

Application	IHC-P, E
Primary Accession	Q12866
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	110kDa KDa
Dilution	IHC-P~~N/A E~~N/A

MER / MERTK Antibody (C-Terminus) - Additional Information**Gene ID** 10461**Other Names**

Tyrosine-protein kinase Mer, 2.7.10.1, Proto-oncogene c-Mer, Receptor tyrosine kinase MerTK, MERTK, MER

Target/Specificity

Human MERTK.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

MER / MERTK Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

MER / MERTK Antibody (C-Terminus) - Protein Information**Name** MERTK**Synonyms** MER**Function**

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to several ligands including LGALS3, TUB, TULP1 or GAS6. Regulates many physiological processes including cell survival, migration, differentiation, and phagocytosis of apoptotic cells (efferocytosis). Ligand binding at the cell surface induces autophosphorylation of MERTK on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with GRB2 or PLCG2 and induces phosphorylation of MAPK1, MAPK2, FAK/PTK2 or RAC1. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and

engulfment (PubMed:32640697). Functions in the retinal pigment epithelium (RPE) as a regulator of rod outer segments fragments phagocytosis. Also plays an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines. Highly expressed in testis, ovary, prostate, lung, and kidney, with lower expression in spleen, small intestine, colon, and liver

MER / MERTK Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MER / MERTK Antibody (C-Terminus) - Images



Anti-MERTK antibody IHC of human adrenal.

MER / MERTK Antibody (C-Terminus) - Background

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to several ligands including LGALS3, TUB, TULP1 or GAS6. Regulates many physiological processes including cell survival, migration, differentiation, and phagocytosis of apoptotic cells (efferocytosis). Ligand binding at the cell surface induces autophosphorylation of MERTK on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with GRB2 or PLCG2 and induces phosphorylation of MAPK1, MAPK2, FAK/PTK2 or RAC1. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and engulfment.

Functions in the retinal pigment epithelium (RPE) as a regulator of rod outer segments fragments phagocytosis. Plays also an important role in inhibition of Toll- like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3.

MER / MERTK Antibody (C-Terminus) - References

Graham D.K.,et al.Cell Growth Differ. 5:647-657(1994).
Graham D.K.,et al.Cell Growth Differ. 5:1022-1022(1994).
Gal A.,et al.Nat. Genet. 26:270-271(2000).
Hillier L.W.,et al.Nature 434:724-731(2005).
Ling L.,et al.J. Biol. Chem. 271:18355-18362(1996).