

BIRC3 / cIAP2 Antibody
Rabbit Polyclonal Antibody
Catalog # ALS13666**Specification**

BIRC3 / cIAP2 Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q13489
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A

BIRC3 / cIAP2 Antibody - Additional Information**Gene ID** 330**Other Names**

Baculoviral IAP repeat-containing protein 3, 6.3.2.-, Apoptosis inhibitor 2, API2, C-IAP2, IAP homolog C, Inhibitor of apoptosis protein 1, IAP-1, hIAP-1, hIAP1, RING finger protein 49, TNFR2-TRAF-signaling complex protein 1, BIRC3, API2, IAP1, MIHC, RNF49

Target/Specificity

Human cIAP2. Predicted cross-reactivity based on amino acid sequence homology: bovine (84%).

Reconstitution & Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Precautions

BIRC3 / cIAP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BIRC3 / cIAP2 Antibody - Protein Information**Name** BIRC3**Synonyms** API2, MIHC, RNF49**Function**

Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and regulates both canonical and non- canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. The target proteins for its E3 ubiquitin-protein ligase activity include: RIPK1, RIPK2, RIPK3, RIPK4, CASP3, CASP7, CASP8, IKBKE, TRAF1, and

BCL10. Acts as an important regulator of innate immune signaling via regulation of Toll-like receptors (TLRs), Nodlike receptors (NLRs) and RIG-I like receptors (RLRs), collectively referred to as pattern recognition receptors (PRRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase-independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8.

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Highly expressed in fetal lung, and kidney. In the adult, expression is mainly seen in lymphoid tissues, including spleen, thymus and peripheral blood lymphocytes


Volume

50 μ l


BIRC3 / cIAP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BIRC3 / cIAP2 Antibody - Images

Sample (30 ug of whole cell lysate).



Anti-clAP2 antibody IHC of human testis.



Anti-clAP2 antibody IHC of human tonsil.

BIRC3 / clAP2 Antibody - Background

Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin- protein ligase regulating NF-kappa-B signaling and regulates both canonical and non-canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. The target proteins for its E3 ubiquitin- protein ligase activity include: RIPK1, RIPK2, RIPK3, RIPK4, CASP3, CASP7, CASP8, IKBKE, TRAF1, and BCL10. Acts as an important regulator of innate immune signaling via regulation of Toll-like receptors (TLRs), Nodlike receptors (NLRs) and RIG-I like receptors (RLRs), collectively referred to as pattern recognition receptors (PRRs). Protects cells from spontaneous formation of the ripoptosome, a large multi-protein complex that has the capability to kill cancer cells in a caspase-dependent and caspase- independent manner. Suppresses ripoptosome formation by ubiquitinating RIPK1 and CASP8.

BIRC3 / clAP2 Antibody - References

- Rothe M.,et al.Cell 83:1243-1252(1995).
- Liston P.,et al.Nature 379:349-353(1996).
- Uren A.G.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:4974-4978(1996).
- Horrevoets A.J.G.,et al.Blood 93:3418-3431(1999).
- Baens M.,et al.Genes Chromosomes Cancer 29:281-291(2000).