

DPP9 Antibody (clone 6A12)
Mouse Monoclonal Antibody
Catalog # ALS16278**Specification****DPP9 Antibody (clone 6A12) - Product Information**

Application	WB, IHC-P, FC
Primary Accession	Q86T12
Reactivity	Human, Dog
Host	Mouse
Clonality	Monoclonal
Calculated MW	98kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50

DPP9 Antibody (clone 6A12) - Additional Information**Gene ID** 91039**Other Names**

Dipeptidyl peptidase 9, DP9, 3.4.14.5, Dipeptidyl peptidase IV-related protein 2, DPRP-2, Dipeptidyl peptidase IX, DPP IX, Dipeptidyl peptidase-like protein 9, DPLP9, DPP9, DPRP2

Target/Specificity

Human DPP9

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

DPP9 Antibody (clone 6A12) is for research use only and not for use in diagnostic or therapeutic procedures.

DPP9 Antibody (clone 6A12) - Protein Information**Name** DPP9 {ECO:0000303|PubMed:12459266, ECO:0000312|HGNC:HGNC:18648}**Function**

Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2 (PubMed:12662155, PubMed:16475979, PubMed:19667070, PubMed:29382749, PubMed:30291141, PubMed:33731929, PubMed:36112693). Acts as a key inhibitor of caspase-1-dependent monocyte and

macrophage pyroptosis in resting cells by preventing activation of NLRP1 and CARD8 (PubMed:27820798, PubMed:29967349, PubMed:30291141, PubMed:31525884, PubMed:32796818, PubMed:36112693, PubMed:36357533). Sequesters the cleaved C-terminal part of NLRP1 and CARD8, which respectively constitute the active part of the NLRP1 and CARD8 inflammasomes, in a ternary complex, thereby preventing their oligomerization and activation (PubMed:33731929, PubMed:33731932, PubMed:34019797). The dipeptidyl peptidase activity is required to suppress NLRP1 and CARD8; however, neither NLRP1 nor CARD8 are bona fide substrates of DPP9, suggesting the existence of substrate(s) required for NLRP1 and CARD8 inhibition (PubMed:33731929).

Cellular Location

[Isoform 1]: Cytoplasm, cytosol

Tissue Location

Ubiquitously expressed, with highest levels in liver, heart and muscle, and lowest levels in brain

Volume


50 µl

DPP9 Antibody (clone 6A12) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPP9 Antibody (clone 6A12) - Images



Western blot of extracts (35 ug) from 9 different cell lines by using anti-DPP9 monoclonal

antibody.



Anti-DPP9 antibody IHC staining of human kidney.

DPP9 Antibody (clone 6A12) - Background

Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2.

DPP9 Antibody (clone 6A12) - References

- Olsen C.,et al.Gene 299:185-193(2002).
- Qi S.Y.,et al.Biochem. J. 373:179-189(2003).
- Ajami K.,et al.Biochim. Biophys. Acta 1679:18-28(2004).
- Bjelke J.R.,et al.Biochem. J. 396:391-399(2006).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).