

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) Goat Polyclonal Antibody Catalog # ALS16651

Specification

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW Dilution WB, IHC-P, IF <u>P17302</u> <u>2697</u> Human, Mouse, Rat, Monkey, Dog Goat Polyclonal IgG 43008 WB~~1:1000 IHC-P~~N/A IF~~1:50~200

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Additional Information

Gene ID 2697

Other Names GJA1, Connexin-43, DFNB38, Connexin 43, Gap junction alpha-1 protein, GJAL, HSS, Gap junction protein alpha 1, ODDD, ODD, AVSD3, CX43, HLHS1, ODOD, SDTY3

Target/Specificity Detects endogenous levels of total connexin 43.

Reconstitution & Storage PBS, 20% glycerol, 0.05% sodium azide. Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Precautions GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Protein Information

Name GJA1

Synonyms GJAL

Function

Gap junction protein that acts as a regulator of bladder capacity. A gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a critical role in the physiology of hearing by participating in the recycling of potassium to the cochlear endolymph.



Negative regulator of bladder functional capacity: acts by enhancing intercellular electrical and chemical transmission, thus sensitizing bladder muscles to cholinergic neural stimuli and causing them to contract (By similarity). May play a role in cell growth inhibition through the regulation of NOV expression and localization. Plays an essential role in gap junction communication in the ventricles (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction, gap junction. Endoplasmic reticulum {ECO:0000250|UniProtKB:P23242}. Note=Localizes at the intercalated disk (ICD) in cardiomyocytes and the proper localization at ICD is dependent on TMEM65. {ECO:0000250|UniProtKB:P23242}

Tissue Location

Expressed at intercalated disks in the heart (at protein level) (PubMed:11741837, PubMed:18662195). Expressed in the fetal cochlea (PubMed:11741837).


Volume 100 μl

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- <u>Blocking Peptides</u>
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Images



Anti-GJA1 / CX43 / Connexin 43 antibody IHC staining of human uterus, endometrium.





Immunofluorescence. Immunostaining of primary RPE cells with CX43 antibody at 1:100 dilution.



Western blot.

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - Background

Gap junction protein that acts as a regulator of bladder capacity. A gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a critical role in the physiology of hearing by participating in the recycling of potassium to the cochlear endolymph. Negative regulator of bladder functional capacity: acts by enhancing intercellular electrical and chemical transmission, thus sensitizing bladder muscles to cholinergic neural stimuli and causing them to contract (By similarity).

GJA1 / CX43 / Connexin 43 Antibody (C-Terminus) - References

Fishman G.I., et al.J. Cell Biol. 111:589-598(1990). Fishman G.I., et al.Genomics 10:250-256(1991). Haefliger J.-A., et al.Eur. Heart J. 20:1843-1843(1999). Halleck A., et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Ota T., et al.Nat. Genet. 36:40-45(2004).