

CD4 Antibody (clone 5D9)
Mouse Monoclonal Antibody
Catalog # ALS16661**Specification**

CD4 Antibody (clone 5D9) - Product Information

Application	WB, IHC-P
Primary Accession	P01730
Other Accession	920
Reactivity	Human, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Calculated MW	51111
Dilution	WB~~1:1000 IHC-P~~N/A

CD4 Antibody (clone 5D9) - Additional Information**Gene ID** 920**Other Names**

CD4, CD4 antigen, CD4 receptor, CD4mut, CD4 antigen (p55), CD4 molecule

Target/Specificity

Human CD4

Reconstitution & Storage

PBS, pH 7.3, 1% BSA, 50% glycerol, 0.02% sodium azide. Store at -20°C. Minimize freezing and thawing.

Precautions

CD4 Antibody (clone 5D9) is for research use only and not for use in diagnostic or therapeutic procedures.

CD4 Antibody (clone 5D9) - Protein Information**Name** CD4**Function**

Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion

and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T- helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Note=Localizes to lipid rafts (PubMed:12517957, PubMed:9168119). Removed from plasma membrane by HIV- 1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation. Cell surface expression is also down-modulated by HIV-1 Envelope glycoprotein gp160 that interacts with, and sequesters CD4 in the endoplasmic reticulum

Tissue Location


Highly expressed in T-helper cells. The presence of CD4 is a hallmark of T-helper cells which are specialized in the activation and growth of cytotoxic T-cells, regulation of B cells, or activation of phagocytes. CD4 is also present in other immune cells such as macrophages, dendritic cells or NK cells

CD4 Antibody (clone 5D9) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CD4 Antibody (clone 5D9) - Images



Anti-CD4 antibody IHC staining of human spleen.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD4...



Western blot of extracts (35 ug) from 9 different cell lines by using anti-CD4 monoclonal...

CD4 Antibody (clone 5D9) - Background

Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell activation. Induces the aggregation of lipid rafts.

CD4 Antibody (clone 5D9) - References

- Maddon P.J.,et al.Cell 42:93-104(1985).
- Littman D.R.,et al.Cell 55:541-541(1988).
- Ansari-Lari M.A.,et al.Genome Res. 6:314-326(1996).
- Ansari-Lari M.A.,et al.Genome Res. 7:268-280(1997).
- Hodge T.W.,et al.Hum. Immunol. 30:99-104(1991).