

### **GPI Antibody (clone 4B11)**

Mouse Monoclonal Antibody Catalog # ALS16950

### **Specification**

# GPI Antibody (clone 4B11) - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype

Calculated MW

Dilution

63147 WB~~1:1000 IHC-P~~N/A IF~~1:50~200

WB, IHC-P, IF

P06744

Human

Mouse

**Monoclonal** 

2821

IgG1

### GPI Antibody (clone 4B11) - Additional Information

#### **Gene ID 2821**

#### **Other Names**

GPI, AMF, Autocrine motility factor, Glucose phosphate isomerase, Hexose monophosphate isomerase, Hexosephosphate isomerase, Neuroleukin, Oxoisomerase, Phosphoglucose isomerase, Phosphohexomerase, Phosphohexose isomerase, SA36, PGI, SA-36, Glucose-6- ...

#### **Target/Specificity**

Human GPI

#### **Reconstitution & Storage**

PBS, pH 7.3, 1% BSA, 50% glycerol, 0.02% sodium azide. Store at -20°C. Minimize freezing and thawing.

#### **Precautions**

GPI Antibody (clone 4B11) is for research use only and not for use in diagnostic or therapeutic procedures.

### **GPI Antibody (clone 4B11) - Protein Information**

Name GPI {ECO:0000303|PubMed:2387591, ECO:0000312|HGNC:HGNC:4458}

#### **Function**

In the cytoplasm, catalyzes the conversion of glucose-6- phosphate to fructose-6-phosphate, the second step in glycolysis, and the reverse reaction during gluconeogenesis (PubMed:<a href="http://www.uniprot.org/citations/28803808" target="\_blank">28803808</a>). Besides it's role as a glycolytic enzyme, also acts as a secreted cytokine: acts as an angiogenic factor (AMF) that stimulates endothelial cell motility (PubMed:<a



href="http://www.uniprot.org/citations/11437381" target="\_blank">11437381</a>). Acts as a neurotrophic factor, neuroleukin, for spinal and sensory neurons (PubMed:<a href="http://www.uniprot.org/citations/11004567" target="\_blank">11004567</a>, PubMed:<a href="http://www.uniprot.org/citations/3352745" target="\_blank">3352745</a>). It is secreted by lectin-stimulated T-cells and induces immunoglobulin secretion (PubMed:<a href="http://www.uniprot.org/citations/11004567" target="\_blank">11004567</a>, PubMed:<a href="http://www.uniprot.org/citations/3352745" target="\_blank">3352745</a>).


Cellular Location Cytoplasm. Secreted

### **GPI Antibody (clone 4B11) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **GPI Antibody (clone 4B11) - Images**




Anti-GPI antibody IHC staining of human kidney.



Anti-GPI mouse monoclonal antibody immunofluorescent staining of COS7 cells transiently...





# GPI Antibody (clone 4B11) - Background

Besides it's role as a glycolytic enzyme, mammalian GPI can function as a tumor-secreted cytokine and an angiogenic factor (AMF) that stimulates endothelial cell motility. GPI is also a neurotrophic factor (Neuroleukin) for spinal and sensory neurons.

# **GPI Antibody (clone 4B11) - References**

Gurney M.E.,et al.Submitted (MAR-1987) to the EMBL/GenBank/DDBJ databases. Yakirevich E.,et al.Biol. Reprod. 62:1016-1023(2000). Ota T.,et al.Nat. Genet. 36:40-45(2004). Grimwood J.,et al.Nature 428:529-535(2004). Walker J.I.H.,et al.Genomics 7:638-643(1990).