

Anti-XBP1 Antibody (clone 2D9)
Mouse Anti Human Monoclonal Antibody
Catalog # ALS17923**Specification**

Anti-XBP1 Antibody (clone 2D9) - Product Information

Application	WB, IHC-P, E
Primary Accession	P17861
Predicted	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a,k
Calculated MW	28695
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

Anti-XBP1 Antibody (clone 2D9) - Additional Information**Gene ID** 7494**Alias Symbol** **XBP1**
Other Names
XBP1, X-box binding protein 1, X-box-binding protein 1, XBP2, TREB5, XBP-1**Target/Specificity**
Human XBP1**Reconstitution & Storage**
Protein A purified**Precautions**
Anti-XBP1 Antibody (clone 2D9) is for research use only and not for use in diagnostic or therapeutic procedures.**Anti-XBP1 Antibody (clone 2D9) - Protein Information****Name** XBP1 ([HGNC:12801](#))**Function**
Functions as a transcription factor during endoplasmic reticulum (ER) stress by regulating the unfolded protein response (UPR). Required for cardiac myogenesis and hepatogenesis during embryonic development, and the development of secretory tissues such as exocrine pancreas and salivary gland (By similarity). Involved in terminal differentiation of B lymphocytes to plasma cells and production of immunoglobulins (PubMed:11460154). Modulates the cellular response to ER stress in a PIK3R-dependent manner (PubMed:20348923). Binds to the cis-acting X box present in the promoter regions of

major histocompatibility complex class II genes (PubMed:8349596). Involved in VEGF-induced endothelial cell (EC) proliferation and retinal blood vessel formation during embryonic development but also for angiogenesis in adult tissues under ischemic conditions. Also functions as a major regulator of the UPR in obesity-induced insulin resistance and type 2 diabetes for the management of obesity and diabetes prevention (By similarity).

Cellular Location

Endoplasmic reticulum. Note=Colocalizes with ERN1 and KDR in the endoplasmic reticulum in endothelial cells in a vascular endothelial growth factor (VEGF)-dependent manner (PubMed:23529610) [Isoform 2]: Nucleus. Cytoplasm {ECO:0000250|UniProtKB:O35426}. Note=Localizes predominantly in the nucleus. Colocalizes in the nucleus with SIRT1. Translocates into the nucleus in a PIK3R-, ER stress-induced- and/or insulin-dependent manner (By similarity). {ECO:0000250|UniProtKB:O35426}

Tissue Location

Expressed in plasma cells in rheumatoid synovium (PubMed:11460154). Over-expressed in primary breast cancer and metastatic breast cancer cells (PubMed:25280941). Isoform 1 and isoform 2 are expressed at higher level in proliferating as compared to confluent quiescent endothelial cells (PubMed:19416856)

Anti-XBP1 Antibody (clone 2D9) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-XBP1 Antibody (clone 2D9) - Images