

Anti-CDC42 Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18161**Specification**

Anti-CDC42 Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P60953
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	21259
Dilution	WB~~1:1000 IHC-P~~N/A

Anti-CDC42 Antibody - Additional Information**Gene ID 998**

Alias Symbol	CDC42
--------------	-------

Other Names
CDC42, G25K, G25K GTP-binding protein, Growth-regulating protein, CDC42Hs, GTP-binding protein, 25kD

Target/Specificity

Human CDC42

Reconstitution & Storage

Affinity purified

Precautions

Anti-CDC42 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CDC42 Antibody - Protein Information**Name** CDC42 ([HGNC:1736](#))**Function**

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase (PubMed:15642749). Regulates cell migration (PubMed:17038317, PubMed:22843693). In neurons, plays a role in the extension and maintenance of

the formation of filopodia, thin and actin-rich surface projections (PubMed:14978216). Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons. In podocytes, facilitates filopodia and podosomes formation upon DOCK11-activation (PubMed:33523862). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (PubMed:26465210). Upon activation by PLEKHG4B, involved in actin cytoskeletal remodeling during epithelial cell-cell junction formation (PubMed:33310911).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody Cell projection, dendrite {ECO:0000250|UniProtKB:P60766} Note=Localizes to spindle during prometaphase cells. Moves to the central spindle as cells progressed through anaphase to telophase (PubMed:15642749). Localizes at the end of cytokinesis in the intercellular bridge formed between two daughter cells (PubMed:15642749). Its localization is regulated by the activities of guanine nucleotide exchange factor ECT2 and GTPase activating protein RACGAP1 (PubMed:15642749). Colocalizes with NEK6 in the centrosome (PubMed:20873783). In its active GTP-bound form localizes to the leading edge membrane of migrating dendritic cells (By similarity) {ECO:0000250|UniProtKB:P60766, ECO:0000269|PubMed:15642749, ECO:0000269|PubMed:20873783}

Anti-CDC42 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CDC42 Antibody - Images