

**INS Antibody (Ascites)**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM1985a****Specification**

---

**INS Antibody (Ascites) - Product Information**

|                   |  |
|-------------------|--|
| Application       | WB,E   |
| Primary Accession | <a href="#">P01308</a>   |
| Other Accession   | <a href="#">NP_001172027.1</a> , <a href="#">NP_000198.1</a> ,<br><a href="#">NP_001172026.1</a> |
| Reactivity        | Human  |
| Host              | Mouse  |
| Clonality         | Monoclonal   |
| Isotype           | IgM  |
| Calculated MW     | 11981  |
| Antigen Region    | 35-64  |

**INS Antibody (Ascites) - Additional Information****Gene ID** 3630**Other Names**

Insulin, Insulin B chain, Insulin A chain, INS

**Target/Specificity**

This INS antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 35-64 amino acids from human INS.

**Dilution**

WB~~1:1000~8000

E~~Use at an assay dependent concentration.

**Format**

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

INS Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

**INS Antibody (Ascites) - Protein Information****Name** INS**Function** Insulin decreases blood glucose concentration. It increases cell permeability to

monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver.

#### **Cellular Location**


Secreted.

### **INS Antibody (Ascites) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **INS Antibody (Ascites) - Images**



INS Antibody (Cat. #AM1985a) western blot analysis in CEM cell line lysates (35µg/lane). This demonstrates the INS antibody detected the INS protein (arrow).

### **INS Antibody (Ascites) - Background**

After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq].

### **INS Antibody (Ascites) - References**

Hinks, A., et al. Ann. Rheum. Dis. 69(12):2169-2172(2010)

Breuer, T.G., et al. Eur. J. Endocrinol. 163(4):551-558(2010)  
Andersen, M.K., et al. Diabetes Care 33(9):2062-2064(2010)  
Ferron, M., et al. Cell 142(2):296-308(2010)  
Authier, F., et al. J. Biol. Chem. 277(11):9437-9446(2002)