

PSMA5 Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2037b**Specification**

PSMA5 Antibody - Product Information

Application	WB, IF, IHC-P,E
Primary Accession	P28066
Other Accession	O9Z2U1 , Q5E987 , NP_002781.2
Reactivity	Human, Mouse, Rat
Predicted	Bovine
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

PSMA5 Antibody - Additional Information**Gene ID** 5686**Other Names**

Proteasome subunit alpha type-5, Macropain zeta chain, Multicatalytic endopeptidase complex zeta chain, Proteasome zeta chain, PSMA5

Target/Specificity

Purified His-tagged PSMA5 protein(Fragment) was used to produced this monoclonal antibody.

Dilution

WB~~1:1000

IF~~1:25

IHC-P~~1:25

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PSMA5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PSMA5 Antibody - Protein Information**Name** PSMA5 ([HGNC:9534](#))**Function** Component of the 20S core proteasome complex involved in the proteolytic degradation

of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP- dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin- independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).

Cellular Location

Cytoplasm. Nucleus. Note=Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9

Tissue Location


Expressed in fetal brain (at protein level).

PSMA5 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PSMA5 Antibody - Images




Fluorescent image of HeLa cells stained with XAF1 PSMA5 Antibody (Cat#AM2037b). AM2037b was diluted at 1:25 dilution. An Alexa Fluor® 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



PSMA5 Antibody (Cat. #AM2037b) western blot analysis in K562 cell line lysates (35µg/lane). This demonstrates the PSMA5 antibody detected the PSMA5 protein (arrow).



All lanes : Anti-PSMA5 Antibody at 1:1000 dilution Lane 1: human liver lysate Lane 2: Hela whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: Neuro-2a whole cell lysate Lane 6: PC-12 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded H. testis section using PSMA5 Antibody(Cat#AM2037b). AM2037b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

PSMA5 Antibody - Background

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit.

PSMA5 Antibody - References

Kottgen, A., et al. Nat. Genet. 42(5):376-384(2010)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)
Hirano, Y., et al. Nature 437(7063):1381-1385(2005)