

FH Antibody (N-term) (Ascites)
Mouse Monoclonal Antibody (Mab)
Catalog # AM2120a**Specification**

FH Antibody (N-term) (Ascites) - Product Information

Application	WB,E
Primary Accession	P07954
Other Accession	P14408 , P10173 , Q60HF9 , NP_000134.2
Reactivity	Human
Predicted	Monkey, Pig, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Calculated MW	54637
Antigen Region	35-63

FH Antibody (N-term) (Ascites) - Additional Information**Gene ID** 2271**Other Names**

Fumarate hydratase, mitochondrial, Fumarase, FH

Target/Specificity

This FH antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 35-63 amino acids from the N-terminal region of human FH.

Dilution

WB~~1:100~1600

E~~Use at an assay dependent concentration.

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FH Antibody (N-term) (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

FH Antibody (N-term) (Ascites) - Protein Information**Name** FH {ECO:0000303|PubMed:27037871, ECO:0000312|HGNC:HGNC:3700}**Function** Catalyzes the reversible stereospecific interconversion of fumarate to L-malate

(PubMed:[30761759](#)). Experiments in other species have demonstrated that specific isoforms of this protein act in defined pathways and favor one direction over the other (Probable).

Cellular Location

[Isoform Mitochondrial]: Mitochondrion

Tissue Location


Expressed in red blood cells; underexpressed in red blood cells (cytoplasm) of patients with hereditary non-spherocytic hemolytic anemia of unknown etiology.

FH Antibody (N-term) (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FH Antibody (N-term) (Ascites) - Images



FH Antibody (N-term)(Ascites)(Cat. #AM2120a) western blot analysis in Jurkat cell line lysates (35µg/lane). This demonstrates the FH antibody detected the FH protein (arrow).

FH Antibody (N-term) (Ascites) - Background

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy.

FH Antibody (N-term) (Ascites) - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
Allegri, G., et al. J. Inherit. Metab. Dis. 33(4):411-419(2010)
Yogev, O., et al. PLoS Biol. 8 (3), E1000328 (2010) :
Yang, Y., et al. Cancer Genet. Cytogenet. 196(1):45-55(2010)
Rikova, K., et al. Cell 131(6):1190-1203(2007)