

GATA3 Antibody (Center)
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM2264b

Specification

GATA3 Antibody (Center) - Product Information

Application	WB, IHC, IF, FC,E
Primary Accession	P23771
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b, κ
Antigen Region	1-444

GATA3 Antibody (Center) - Additional Information

Gene ID 2625

Other Names

Trans-acting T-cell-specific transcription factor GATA-3, GATA-binding factor 3, GATA3

Target/Specificity

This GATA3 antibody is generated from a mouse immunized with a recombinant protein from human GATA3.

Dilution

WB~~1:1000

IHC~~1:400

IF~~1:25

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GATA3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GATA3 Antibody (Center) - Protein Information

Name GATA3

Function Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses. Positively regulates ASB2 expression (By similarity). Coordinates macrophage transcriptional activation and UCP2-dependent metabolic reprogramming in response to IL33. Upon tissue injury, acts downstream of IL33 signaling to drive differentiation of inflammation-resolving alternatively activated macrophages.

Cellular Location

Nucleus.

Tissue Location


T-cells and endothelial cells.

GATA3 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


GATA3 Antibody (Center) - Images




Fluorescent image of MCF-7 cells stained with GATA3 Antibody (Center)(Cat#AM2264b). AM2264b was diluted at 1:25 dilution. An Alexa Fluor® 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western blot analysis of lysates from CEM, HL-60, SH-SY5Y cell lines (from left to right) using GATA3 Antibody (Center) (Cat. # AM2264b). AM2264b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.



Immunohistochemical analysis of paraffin-embedded Human Normal breast section using Pink1(Cat#AM2264b). AM2264b was diluted at 1:400 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Flow cytometric analysis of MCF-7 cells using GATA3 Antibody (Center)(green, Cat#AM2264b) compared to an isotype control of mouse IgG2b(blue). AM2264b was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.

GATA3 Antibody (Center) - Background

Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'.

GATA3 Antibody (Center) - References

Joulin V.,et al.EMBO J. 10:1809-1816(1991).
Ho I.-C.,et al.EMBO J. 10:1187-1192(1991).
Ko L.J.,et al.Mol. Cell. Biol. 11:2778-2784(1991).
Marine J.,et al.Proc. Natl. Acad. Sci. U.S.A. 88:7284-7288(1991).
Deloukas P.,et al.Nature 429:375-381(2004).