

SQSTM1 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8447b

Specification

SQSTM1 Antibody - Product Information

Application	WB,E
Primary Accession	O13501
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,κ

SQSTM1 Antibody - Additional Information

Gene ID 8878

Other Names

Sequestosome-1, EBI3-associated protein of 60 kDa, EBIAP, p60, Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa, Ubiquitin-binding protein p62, SQSTM1, ORCA, OSIL

Target/Specificity

This SQSTM1 antibody is generated from mice immunized with a recombinant protein.

Dilution

WB~~1:2000
E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SQSTM1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SQSTM1 Antibody - Protein Information

Name SQSTM1 {ECO:0000303|PubMed:16286508, ECO:0000312|HGNC:HGNC:11280}

Function Molecular adapter required for selective macroautophagy (aggrephagy) by acting as a bridge between polyubiquitinated proteins and autophagosomes (PubMed:[15340068](#), PubMed:[15953362](#), PubMed:[16286508](#), PubMed:[17580304](#), PubMed:[20168092](#), PubMed:[22017874](#), PubMed:[22622177](#), PubMed:[24128730](#), PubMed:[28404643](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[31857589](#), PubMed:[33509017](#),

PubMed:[34471133](#), PubMed:[34893540](#), PubMed:[35831301](#), PubMed:[37306101](#), PubMed:[37802024](#)). Promotes the recruitment of ubiquitinated cargo proteins to autophagosomes via multiple domains that bridge proteins and organelles in different steps (PubMed:[16286508](#), PubMed:[20168092](#), PubMed:[22622177](#), PubMed:[24128730](#), PubMed:[28404643](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[34893540](#), PubMed:[37802024](#)). SQSTM1 first mediates the assembly and removal of ubiquitinated proteins by undergoing liquid-liquid phase separation upon binding to ubiquitinated proteins via its UBA domain, leading to the formation of insoluble cytoplasmic inclusions, known as p62 bodies (PubMed:[15911346](#), PubMed:[20168092](#), PubMed:[22017874](#), PubMed:[24128730](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[31857589](#), PubMed:[37802024](#)). SQSTM1 then interacts with ATG8 family proteins on autophagosomes via its LIR motif, leading to p62 body recruitment to autophagosomes, followed by autophagic clearance of ubiquitinated proteins (PubMed:[16286508](#), PubMed:[17580304](#), PubMed:[20168092](#), PubMed:[22622177](#), PubMed:[24128730](#), PubMed:[28404643](#), PubMed:[37802024](#)). SQSTM1 is itself degraded along with its ubiquitinated cargos (PubMed:[16286508](#), PubMed:[17580304](#), PubMed:[37802024](#)). Also required to recruit ubiquitinated proteins to PML bodies in the nucleus (PubMed:[20168092](#)). Also involved in autophagy of peroxisomes (pexophagy) in response to reactive oxygen species (ROS) by acting as a bridge between ubiquitinated PEX5 receptor and autophagosomes (PubMed:[26344566](#)). Acts as an activator of the NFE2L2/NRF2 pathway via interaction with KEAP1: interaction inactivates the BCR(KEAP1) complex by sequestering the complex in inclusion bodies, promoting nuclear accumulation of NFE2L2/NRF2 and subsequent expression of cytoprotective genes (PubMed:[20452972](#), PubMed:[28380357](#), PubMed:[33393215](#), PubMed:[37306101](#)). Promotes relocalization of 'Lys-63'-linked ubiquitinated STING1 to autophagosomes (PubMed:[29496741](#)). Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (PubMed:[27368102](#), PubMed:[33472082](#)). Sequesters tensin TNS2 into cytoplasmic puncta, promoting TNS2 ubiquitination and proteasomal degradation (PubMed:[25101860](#)). May regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1 (PubMed:[10356400](#), PubMed:[10747026](#), PubMed:[11244088](#), PubMed:[12471037](#), PubMed:[16079148](#), PubMed:[19931284](#)). May play a role in titin/TTN downstream signaling in muscle cells (PubMed:[15802564](#)). Adapter that mediates the interaction between TRAF6 and CYLD (By similarity).

Cellular Location

Cytoplasmic vesicle, autophagosome. Preautophagosomal structure. Cytoplasm, cytosol. Nucleus, PML body. Late endosome. Lysosome. Nucleus Endoplasmic reticulum. Cytoplasm, myofibril, sarcomere {ECO:0000250|UniProtKB:O08623}. Note=In cardiac muscle, localizes to the sarcomeric band (By similarity). Localizes to cytoplasmic membraneless inclusion bodies, known as p62 bodies, containing polyubiquitinated protein aggregates (PubMed:[11786419](#), PubMed:[20357094](#), PubMed:[22017874](#), PubMed:[29343546](#), PubMed:[29507397](#), PubMed:[31857589](#), PubMed:[37306101](#), PubMed:[37802024](#)). In neurodegenerative diseases, detected in Lewy bodies in Parkinson disease, neurofibrillary tangles in Alzheimer disease, and HTT aggregates in Huntington disease (PubMed:[15158159](#)). In protein aggregate diseases of the liver, found in large amounts in Mallory bodies of alcoholic and nonalcoholic steatohepatitis, hyaline bodies in hepatocellular carcinoma, and in SERPINA1 aggregates (PubMed:[11981755](#)) Enriched in Rosenthal fibers of pilocytic astrocytoma (PubMed:[11786419](#)). In the cytoplasm, observed in both membrane-free ubiquitin-containing protein aggregates (sequestosomes) and membrane- surrounded autophagosomes (PubMed:[15953362](#), PubMed:[17580304](#)) Colocalizes with TRIM13 in the perinuclear endoplasmic reticulum (PubMed:[22178386](#)). Co-localizes with TRIM5 in cytoplasmic bodies (PubMed:[20357094](#)). When nuclear export is blocked by treatment with leptomycin B, accumulates in PML bodies (PubMed:[20168092](#)) {ECO:0000250|UniProtKB:O08623, ECO:0000269|PubMed:[11786419](#), ECO:0000269|PubMed:[11981755](#), ECO:0000269|PubMed:[15158159](#), ECO:0000269|PubMed:[15953362](#), ECO:0000269|PubMed:[17580304](#), ECO:0000269|PubMed:[20168092](#), ECO:0000269|PubMed:[20357094](#), ECO:0000269|PubMed:[22178386](#), ECO:0000269|PubMed:[29343546](#),

ECO:0000269|PubMed:29507397, ECO:0000269|PubMed:31857589,
ECO:0000269|PubMed:37306101, ECO:0000269|PubMed:37802024}

Tissue Location


Ubiquitously expressed.

SQSTM1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SQSTM1 Antibody - Images



All lanes : Anti-SQSTM1 Antibody at 1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: NCI-H1299 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 62 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SQSTM1 Antibody - Background

Required both for the formation and autophagic degradation of polyubiquitin-containing bodies, called ALIS (aggresome-like induced structures). Links ALIS to the autophagic machinery via direct interaction with MAP1 LC3 family members. May regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. May play a role in titin/TTN downstream signaling in muscle cells. May regulate signaling cascades through ubiquitination. Adapter that mediates the interaction between TRAF6 and CYLD (By similarity). May be involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels.

SQSTM1 Antibody - References

Devergne O.,et al.J. Virol. 70:1143-1153(1996).
Joung I.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:5991-5995(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Schmutz J.,et al.Nature 431:268-274(2004).
Vadlamudi R.K.,et al.FEBS Lett. 435:138-142(1998).