

RAB27B Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AM8508b

Specification

RAB27B Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype

,E <u>000194</u> Human, Mouse Mouse monoclonal IgG1,k

RAB27B Antibody - Additional Information

Gene ID 5874

Other Names Ras-related protein Rab-27B, C25KG, RAB27B

Target/Specificity

This RAB27B antibody is generated from a mouse immunized with a recombinant protein of human RAB27B.

Dilution

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions RAB27B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB27B Antibody - Protein Information

Name RAB27B

Function Small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate homeostasis of late endocytic pathway, including endosomal positioning, maturation and secretion (PubMed:<u>30771381</u>). Plays a role in NTRK2/TRKB axonal anterograde transport by facilitating the association of NTRK2/TRKB with KLC1 (PubMed:<u>21775604</u>). May be involved in targeting uroplakins to urothelial apical membranes (By similarity).



Cellular Location Membrane; Lipid-anchor. Late endosome


Tissue Location Expressed primarily in testis.

RAB27B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

RAB27B Antibody - Images



Anti-RAB27B Antibody at 1:2000 dilution + mouse stomach lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RAB27B Antibody - Background

May be involved in targeting uroplakins to urothelial apical membranes.

RAB27B Antibody - References

Chen D.,et al.Biochem. Mol. Med. 60:27-37(1997). Chen D.,et al.Submitted (JUN-1999) to the EMBL/GenBank/DDBJ databases. Ramalho J.S.,et al.BMC Genet. 2:2-2(2001). Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).