

FZR Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8522b

Specification

FZR Antibody - Product Information

Application WB,E
Primary Accession Q9UM11
Reactivity Human
Host Mouse
Clonality monoclonal
Isotype IgG1,k
Calculated MW 55179

FZR Antibody - Additional Information

Gene ID 51343

Other Names

Fizzy-related protein homolog, Fzr, CDC20-like protein 1, Cdh1/Hct1 homolog, hCDH1, FZR1, CDH1, FYR, FZR, KIAA1242

Target/Specificity

This FZR antibody is generated from a mouse immunized with a recombinant protein between 1-496 amino acids from human FZR.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FZR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FZR Antibody - Protein Information

Name FZR1 (HGNC:24824)

Function Substrate-specific adapter for the anaphase promoting complex/cyclosome (APC/C) E3 ubiquitin-protein ligase complex. Associates with the APC/C in late mitosis, in replacement of CDC20, and activates the APC/C during anaphase and telophase. The APC/C remains active in



Tel: 858.875.1900 Fax: 858.875.1999

degrading substrates to ensure that positive regulators of the cell cycle do not accumulate prematurely. At the G1/S transition FZR1 is phosphorylated, leading to its dissociation from the APC/C. Following DNA damage, it is required for the G2 DNA damage checkpoint: its dephosphorylation and reassociation with the APC/C leads to the ubiquitination of PLK1, preventing entry into mitosis. Acts as an adapter for APC/C to target the DNA-end resection factor RBBP8/CtIP for ubiquitination and subsequent proteasomal degradation. Through the regulation of RBBP8/CtIP protein turnover, may play a role in DNA damage response, favoring DNA double-strand repair through error-prone non-homologous end joining (NHEJ) over error-free, RBBP8-mediated homologous recombination (HR) (PubMed:25349192).

Cellular Location [Isoform 2]: Nucleus

Tissue Location


Isoform 2 is expressed at high levels in heart, liver, spleen and some cancer cell lines whereas isoform 3 is expressed only at low levels in these tissues.

FZR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FZR Antibody - Images



All lanes : Anti-FZR Antibody at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: Hela whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



FZR Antibody - Background

Key regulator of ligase activity of the anaphase promoting complex/cyclosome (APC/C), which confers substrate specificity upon the complex. Associates with the APC/C in late mitosis, in replacement of CDC20, and activates the APC/C during anaphase and telophase. The APC/C remains active in degrading substrates to ensure that positive regulators of the cell cycle do not accumulate prematurely. At the G1/S transition FZR1 is phosphorylated, leading to its dissociation from the APC/C. Following DNA damage, it is required for the G2 DNA damage checkpoint: its dephosphorylation and reassociation with the APC/C leads to the ubiquitination of PLK1, preventing entry into mitosis.

FZR Antibody - References

Kramer E.R.,et al.Curr. Biol. 8:1207-1210(1998). Kotani S.,et al.Submitted (APR-1998) to the EMBL/GenBank/DDBJ databases. Sudo T.,et al.Submitted (JUL-1998) to the EMBL/GenBank/DDBJ databases. Zhou Y.,et al.Biochem. J. 374:349-358(2003). Nagase T.,et al.DNA Res. 6:337-345(1999).