

Anti- β -Catenin (Tyr-333) [γ -Catenin (Tyr-324)], Phosphospecific Antibody
Catalog # AN1678**Specification****Anti- β -Catenin (Tyr-333) [γ -Catenin (Tyr-324)], Phosphospecific Antibody - Product Information**

Primary Accession	P35222
Reactivity	Bovine, Chicken, Drosophila, C.Elegans
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	85497

Anti- β -Catenin (Tyr-333) [γ -Catenin (Tyr-324)], Phosphospecific Antibody - Additional InformationGene ID **1499****Other Names**

Catenin beta1, CTNNB1, catenin

Target/Specificity

β -Catenin is a 92 kDa protein that binds to the cytoplasmic tail of E-Cadherin. The cadherins, transmembrane adhesion molecules, are found with catenins at adherens junctions. Deletions in the cytoplasmic domain of E-Cadherin eliminate catenin binding and result in a loss of cell adhesion. Tyrosine phosphorylation of β -Catenin can regulate its interaction with critical components of adherens junctions. Both Fer and Fyn kinases phosphorylate tyrosine 142 in vitro. Overexpression of these kinases in epithelial cells disrupts interactions between α - and β -Catenins. The phosphorylation of tyrosine 142 may act as a switch from the transcriptional to the adhesive role of β -Catenin. Src family kinases can also phosphorylate tyrosine 86 and 654 in β -Catenin. The Tyr-654 phosphorylation regulates β -Catenin binding to E-cadherin. Thus, site-specific tyrosine phosphorylation of β -Catenin may regulate protein-protein interactions leading to changes in cell adhesion.

Format

Antigen Affinity Purified

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti- β -Catenin (Tyr-333) [γ -Catenin (Tyr-324)], Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Anti- β -Catenin (Tyr-333) [γ -Catenin (Tyr-324)], Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-β-Catenin (Tyr-333) [γ-Catenin (Tyr-324)], Phosphospecific Antibody - Images



Western blot analysis of A431 cells stimulated with pervanadate (1 mM) for 30 min (lanes 1 & 3) then treated with alkaline phosphatase (lanes 2 & 4). The blot was probed with CP1061 anti-β-Catenin (CP1061) (lanes 1 & 2) and CP3051 anti-β-Catenin (Tyr-333) (lanes 3 & 4).

Anti-β-Catenin (Tyr-333) [γ-Catenin (Tyr-324)], Phosphospecific Antibody - Background

β-Catenin is a 92 kDa protein that binds to the cytoplasmic tail of E-Cadherin. The cadherins, transmembrane adhesion molecules, are found with catenins at adherens junctions. Deletions in the cytoplasmic domain of E-Cadherin eliminate catenin binding and result in a loss of cell adhesion. Tyrosine phosphorylation of β-Catenin can regulate its interaction with critical components of adherens junctions. Both Fer and Fyn kinases phosphorylate tyrosine 142 in vitro. Overexpression of these kinases in epithelial cells disrupts interactions between α- and β-Catenins. The phosphorylation of tyrosine 142 may act as a switch from the transcriptional to the adhesive role of β-Catenin. Src family kinases can also phosphorylate tyrosine 86 and 654 in β-Catenin. The Tyr-654 phosphorylation regulates β-Catenin binding to E-cadherin. Thus, site-specific tyrosine phosphorylation of β-Catenin may regulate protein-protein interactions leading to changes in cell adhesion.