

Anti-Prion Protein (Ser-43), Phosphospecific Antibody
Catalog # AN1919**Specification****Anti-Prion Protein (Ser-43), Phosphospecific Antibody - Product Information**

Primary Accession	P04156
Reactivity	Bovine
Host	Rabbit
Clonality	Rabbit Polyclonal
Isotype	IgG
Calculated MW	27661

Anti-Prion Protein (Ser-43), Phosphospecific Antibody - Additional Information

Gene ID	5621
Other Names	
PrP, PrPsc, PrPc	

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Prion Protein (Ser-43), Phosphospecific Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Shipping


Blue Ice

Anti-Prion Protein (Ser-43), Phosphospecific Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Prion Protein (Ser-43), Phosphospecific Antibody - Images



Western blot of GST recombinant human full-length prion protein that was untreated (lanes 1 and 3) or phosphorylated with Cdk5/p25 (lanes 2 & 4). Endogenous prion phosphorylation was examined in human PC3 cells untreated (lanes 5 & 7) or treated with Calyculin A (100 nM) for 30 min (lanes 6 & 8). The blots were probed with anti-Prion protein (3F4) (lanes 1, 2, 5, & 6) or anti-Prion protein (Ser-43) (lanes 3, 4, 7, & 8).

Anti-Prion Protein (Ser-43), Phosphospecific Antibody - Background

Prion related neurodegenerative diseases, called transmissible spongiform encephalopathies, are observed in many animal species. These diseases involve conversion of normal cellular prion protein (PrP^c) into a form that is insoluble and resistant to proteases (PrP^{Sc}). The protease resistant form can polymerize into fibrils which accumulate in infected tissues and cause cell death and tissue damage. PrPs have an N-terminal signal sequence and a C-terminal linkage to glycosylphosphatidylinositol anchor. The mature protein is a glycosylated protein that associates with cell membranes. Phosphorylation of PrP^c at Ser-43 by Cdk5 promotes proteinase K resistance, prion aggregation, and fibril formation in vitro. In addition, Ser-43 phosphorylation is upregulated in scrapie-infected mouse brain relative to controls. Thus, phosphorylation of Ser-43 may be an important mechanism leading conversion of PrP^c to PrP^{Sc} and the onset of disease.