

Anti-Stat3 (N-terminal region) Antibody
Catalog # AN1979**Specification****Anti-Stat3 (N-terminal region) Antibody - Product Information**

Application	WB
Primary Accession	P40763
Reactivity	Bovine
Host	Mouse
Clonality	Mouse Monoclonal
Isotype	IgG1
Calculated MW	88068

Anti-Stat3 (N-terminal region) Antibody - Additional InformationGene ID **6774****Other Names**

Signal transducer and activator of transcription 3

Dilution

WB~~1:1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Stat3 (N-terminal region) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Shipping


Blue Ice

Anti-Stat3 (N-terminal region) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Stat3 (N-terminal region) Antibody - Images



Western blot analysis of human A431 cells untreated (lanes 1, 3, 5, 7, & 9) or treated with EGF (100 nM) for 60 min (lanes 2, 4, 6, 8, & 10). The blots were probed with anti-Stat1 (lanes 1 & 2), anti-Stat1 (Tyr-701) (lanes 3 & 4), anti-Stat3 (lanes 5 & 6), anti-Stat5 (lanes 7 & 8), and anti-Stat5 (Tyr-694) (lanes 9 & 10).

Anti-Stat3 (N-terminal region) Antibody - Background

The stat proteins (Stat1-6) function both as cytoplasmic signal transducers and as activators of transcription in response to cytokines and growth factor receptors. Stat3 is expressed as two variants, Stat3 α (86 kDa) and Stat3 β (79 kDa) that can differ in expression and activity depending on cell type, activation pathway, and cell maturation stage. Both are activated by phosphorylation at Tyr-705, which induces dimerization, nuclear translocation and DNA binding. Stat3 α (86 kDa) transcriptional activation may be regulated by phosphorylation at Ser-727 through the MAPK pathway, while Stat3 β lacks this serine site.