

ATM Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1525a**Specification****ATM Antibody - Product Information**

Application	WB, E
Primary Accession	Q13315
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	351kDa KDa

Description

The protein encoded by this gene belongs to the PI3/PI4-kinase family. This protein is an important cell cycle checkpoint kinase that phosphorylates; thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability. Mutations in this gene are associated with ataxia telangiectasia, an autosomal recessive disorder. Tissue specificity: Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes.

Immunogen

Purified recombinant fragment of human ATM expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

ATM Antibody - Additional Information

Gene ID 472

Other Names

Serine-protein kinase ATM, 2.7.11.1, Ataxia telangiectasia mutated, A-T mutated, ATM

Dilution

WB~~1/500 - 1/2000

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ATM Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

[>26240375, PubMed:26774286, PubMed:30171069, PubMed:30612738, PubMed:30886146, PubMed:30952868, PubMed:38128537, PubMed:9733515, PubMed:9843217\). May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation \(PubMed:19965871\). Phosphorylates ATF2 which stimulates its function in DNA damage response \(PubMed:15916964\). Phosphorylates ERCC6 which is essential for its chromatin remodeling activity at DNA double-strand breaks \(PubMed:29203878\). Phosphorylates TTC5/STRAP at 'Ser-203' in the cytoplasm in response to DNA damage, which promotes TTC5/STRAP nuclear localization \(PubMed:15448695\). Also involved in pexophagy by mediating phosphorylation of PEX5: translocated to peroxisomes in response to reactive oxygen species \(ROS\), and catalyzes phosphorylation of PEX5, promoting PEX5 ubiquitination and induction of pexophagy \(PubMed:26344566\).](http://www.uniprot.org/citations/26240375)

Cellular Location

Nucleus. Cytoplasmic vesicle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q62388}. Peroxisome matrix. Note=Primarily nuclear (PubMed:9050866, PubMed:9150358). Found also in endocytic vesicles in association with beta-adaptin (PubMed:9707615). Translocated to peroxisomes in response to reactive oxygen species (ROS) by PEX5 (PubMed:26344566)

Tissue Location

Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes

ATM Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Figure 1: Western blot analysis using ATM mAb against HEK293 (1) and ATM(AA: 2705-2820)-hlgGfc transfected HEK293 (2) cell lysate.

ATM Antibody - References

1. Am J Hum Genet. 2008 Aug;83(2):243-53.
2. Mol Cell. 2008 Jul 25;31(2):167-77.