

**HIF1A Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1546a****Specification****HIF1A Antibody - Product Information**

Application	WB, IHC, ICC, E
Primary Accession	<a href="#">Q16665</a>
Reactivity	Human, Mouse, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	120kDa KDa

**Description**

Hypoxia-inducible factor-1 (HIF1) is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). This gene encodes the alpha subunit of HIF-1. Overexpression of a natural antisense transcript (aHIF) of this gene has been shown to be associated with nonpapillary renal carcinomas. Two alternative transcripts encoding different isoforms have been identified. (provided by RefSeq)  
Tissue specificity: Expressed in most tissues with highest levels in kidney and heart.  
Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors.

**Immunogen**

Purified recombinant fragment of human HIF1A expressed in E. Coli.

**Formulation**

Ascitic fluid containing 0.03% sodium azide.

**HIF1A Antibody - Additional Information**

**Gene ID** 3091

**Other Names**

Hypoxia-inducible factor 1-alpha, HIF-1-alpha, HIF1-alpha, ARNT-interacting protein, Basic-helix-loop-helix-PAS protein MOP1, Class E basic helix-loop-helix protein 78, bHLHe78, Member of PAS protein 1, PAS domain-containing protein 8, HIF1A, BHLHE78, MOP1, PASD8

**Dilution**

WB~~1/500 - 1/2000  
IHC~~1/200 - 1/1000  
ICC~~N/A  
E~~1/10000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small

aliquots to prevent freeze-thaw cycles.

### Precautions

HIF1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## HIF1A Antibody - Protein Information

**Name** HIF1A {ECO:0000303|PubMed:7539918, ECO:0000312|HGNC:HGNC:4910}

### Function

Functions as a master transcriptional regulator of the adaptive response to hypoxia (PubMed:<a href="http://www.uniprot.org/citations/11292861" target="\_blank">11292861</a>, PubMed:<a href="http://www.uniprot.org/citations/11566883" target="\_blank">11566883</a>, PubMed:<a href="http://www.uniprot.org/citations/15465032" target="\_blank">15465032</a>, PubMed:<a href="http://www.uniprot.org/citations/16973622" target="\_blank">16973622</a>, PubMed:<a href="http://www.uniprot.org/citations/17610843" target="\_blank">17610843</a>, PubMed:<a href="http://www.uniprot.org/citations/18658046" target="\_blank">18658046</a>, PubMed:<a href="http://www.uniprot.org/citations/20624928" target="\_blank">20624928</a>, PubMed:<a href="http://www.uniprot.org/citations/22009797" target="\_blank">22009797</a>, PubMed:<a href="http://www.uniprot.org/citations/30125331" target="\_blank">30125331</a>, PubMed:<a href="http://www.uniprot.org/citations/9887100" target="\_blank">9887100</a>). Under hypoxic conditions, activates the transcription of over 40 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia (PubMed:<a href="http://www.uniprot.org/citations/11292861" target="\_blank">11292861</a>, PubMed:<a href="http://www.uniprot.org/citations/11566883" target="\_blank">11566883</a>, PubMed:<a href="http://www.uniprot.org/citations/15465032" target="\_blank">15465032</a>, PubMed:<a href="http://www.uniprot.org/citations/16973622" target="\_blank">16973622</a>, PubMed:<a href="http://www.uniprot.org/citations/17610843" target="\_blank">17610843</a>, PubMed:<a href="http://www.uniprot.org/citations/20624928" target="\_blank">20624928</a>, PubMed:<a href="http://www.uniprot.org/citations/22009797" target="\_blank">22009797</a>, PubMed:<a href="http://www.uniprot.org/citations/30125331" target="\_blank">30125331</a>, PubMed:<a href="http://www.uniprot.org/citations/9887100" target="\_blank">9887100</a>). Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease (PubMed:<a href="http://www.uniprot.org/citations/22009797" target="\_blank">22009797</a>). Heterodimerizes with ARNT; heterodimer binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters (By similarity). Activation requires recruitment of transcriptional coactivators such as CREBBP and EP300 (PubMed:<a href="http://www.uniprot.org/citations/16543236" target="\_blank">16543236</a>, PubMed:<a href="http://www.uniprot.org/citations/9887100" target="\_blank">9887100</a>). Activity is enhanced by interaction with NCOA1 and/or NCOA2 (PubMed:<a href="http://www.uniprot.org/citations/10594042" target="\_blank">10594042</a>). Interaction with redox regulatory protein APEX1 seems to activate CTAD and potentiates activation by NCOA1 and CREBBP (PubMed:<a href="http://www.uniprot.org/citations/10202154" target="\_blank">10202154</a>, PubMed:<a href="http://www.uniprot.org/citations/10594042" target="\_blank">10594042</a>). Involved in the axonal distribution and transport of mitochondria in neurons during hypoxia (PubMed:<a href="http://www.uniprot.org/citations/19528298" target="\_blank">19528298</a>).

### Cellular Location

Cytoplasm. Nucleus. Nucleus speckle {ECO:0000250|UniProtKB:Q61221}. Note=Colocalizes with HIF3A in the nucleus and speckles (By similarity). Cytoplasmic in normoxia, nuclear translocation in response to hypoxia (PubMed:9822602) {ECO:0000250|UniProtKB:Q61221, ECO:0000269|PubMed:9822602}

### Tissue Location

Expressed in most tissues with highest levels in kidney and heart. Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors. A higher level expression seen in pituitary tumors as compared to the pituitary gland.

### HIF1A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)




Figure 1: Western blot analysis using HIF1A mouse mAb against Cos7 (1), HeLa (2), Jurkat (3), RAJI (4) and NIH/3T3 (5) cell lysate.




Figure 2: Immunohistochemical analysis of paraffin-embedded liver cancer tissues (left) and lung cancer tissues (right) using HIF1A mouse mAb with DAB staining.




Figure 3: Immunohistochemical analysis of paraffin-embedded stomach cancer tissues (left) and brain tumor tissues (right) using HIF1A mouse mAb with DAB staining.




Figure 4: Immunofluorescence analysis of Hela cells using HIF1A mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

#### HIF1A Antibody - References

1. Int J Radiat Oncol Biol Phys. 2008 Dec 1;72(5):1551-9.
2. Eur J Appl Physiol. 2009 Mar;105(4):515-24.