

DLL4 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1723a**Specification****DLL4 Antibody - Product Information**

Application	WB, FC, E
Primary Accession	Q9NR61
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	4A11F8 IgG2b/4A11G2 IgG1
Calculated MW	74.6kDa KDa

Description

This gene is a homolog of the Drosophila delta gene. The delta gene family encodes Notch ligands that are characterized by a DSL domain, EGF repeats, and a transmembrane domain.

Immunogen

Purified recombinant fragment of human DLL4 expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

DLL4 Antibody - Additional Information

Gene ID 54567

Other Names

Delta-like protein 4, Drosophila Delta homolog 4, Delta4, DLL4

Dilution

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DLL4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DLL4 Antibody - Protein Information

Name DLL4

Function

Involved in the Notch signaling pathway as Notch ligand (PubMed:11134954). Activates NOTCH1 and NOTCH4. Involved in angiogenesis; negatively regulates endothelial cell proliferation and migration and angiogenic sprouting (PubMed:20616313). Essential for retinal progenitor proliferation. Required for suppressing rod fates in late retinal progenitors as well as for proper generation of other retinal cell types (By similarity). During spinal cord neurogenesis, inhibits V2a interneuron fate (PubMed:17728344).

Cellular Location

Cell membrane; Single-pass type I membrane protein


Tissue Location

Expressed in vascular endothelium.

DLL4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using DLL4 mAb against human DLL4 (AA: 313-439) recombinant protein. (Expected MW is 39.2 kDa)




Figure 2: Western blot analysis using DLL4 mAb against HEK293 (1) and DLL4 (AA: 313-439)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using DLL4 mouse mAb (green) and negative control (red).

DLL4 Antibody - References

1. Blood. 2010 Sep 30;116(13):2385-94. 2. Circ Res. 2010 Jul 23;107(2):283-93.