

CDKN2A Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1781a**Specification****CDKN2A Antibody - Product Information**

Application	WB, IHC, FC, E
Primary Accession	P42771
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	16.5kDa KDa

Description

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, MDM1, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Immunogen

Purified recombinant fragment of human CDKN2A (AA: 1-156) expressed in E. Coli.

Formulation

Ascitic fluid containing 0.03% sodium azide.

CDKN2A Antibody - Additional Information

Gene ID 1029

Other Names

Cyclin-dependent kinase inhibitor 2A, isoforms 1/2/3, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A, CDKN2, MTS1

Dilution

WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
FC~~1/200 - 1/400
E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small

aliquots to prevent freeze-thaw cycles.

Precautions

CDKN2A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CDKN2A Antibody - Protein Information

Name CDKN2A ([HGNC:1787](#))

Synonyms CDKN2, MTS1

Function

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

Cellular Location

Cytoplasm. Nucleus


Tissue Location

Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific

CDKN2A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CDKN2A mAb against human CDKN2A recombinant protein. (Expected MW is 19 kDa)




Figure 2: Western blot analysis using CDKN2A mAb against HEK293 (1) and CDKN2A (AA: 1-156)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of HEK293 cells using CDKN2A mouse mAb (green) and negative control (red).




Figure 4: Immunohistochemical analysis of paraffin-embedded lung cancer tissues using CDKN2A mouse mAb with DAB staining.




Figure 5: Immunohistochemical analysis of paraffin-embedded endometrial cancer tissues using CDKN2A mouse mAb with DAB staining.

CDKN2A Antibody - References

1.Clin Cancer Res. 2011 Dec 1;17(23):7413-23. 2.Appl Immunohistochem Mol Morphol. 2011 Dec;19(6):562-8.