

MAP2K3 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2234a**Specification****MAP2K3 Antibody - Product Information**

Application	WB, IHC, FC, ICC, E
Primary Accession	P46734
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	39.3kDa KDa

Description

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersinia pseudotuberculosis. Multiple alternatively spliced transcript variants that encode distinct isoforms have been reported for this gene.

Immunogen

Purified recombinant fragment of human MAP2K3 (AA: 1-138) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

MAP2K3 Antibody - Additional Information

Gene ID 5606

Other Names

Dual specificity mitogen-activated protein kinase kinase 3, MAP kinase kinase 3, MAPKK 3, 2.7.12.2, MAPK/ERK kinase 3, MEK 3, Stress-activated protein kinase kinase 2, SAPK kinase 2, SAPKK-2, SAPKK2, MAP2K3, MEK3, MKK3, PRKMK3, SKK2

Dilution

WB~~1/500 - 1/2000
IHC~~1/200 - 1/1000
FC~~1/200 - 1/400
ICC~~N/A
E~~1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MAP2K3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MAP2K3 Antibody - Protein Information

Name MAP2K3

Synonyms MEK3, MKK3, PRKMK3, SKK2

Function

Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14.

Tissue Location

Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues

MAP2K3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)