

## **TGFBR3 Antibody**

Purified Mouse Monoclonal Antibody Catalog # AO2253a

## **Specification**

## **TGFBR3 Antibody - Product Information**

Application WB, E
Primary Accession O03167

Reactivity Human, Mouse

Host Mouse
Clonality Monoclonal
Isotype IgG1

Calculated MW 93.4kDa KDa

**Description** 

This locus encodes the transforming growth factor (TGF)-beta type III receptor. The encoded receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFBR3, which may inhibit TGFB signaling. Decreased expression of this receptor has been observed in various cancers. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

### **Immunogen**

Purified recombinant fragment of human TGFBR3 (AA: 147-328) expressed in E. Coli.

#### Formulation

Purified antibody in PBS with 0.05% sodium azide

## **TGFBR3 Antibody - Additional Information**

## **Gene ID** 7049

### **Other Names**

Transforming growth factor beta receptor type 3, TGF-beta receptor type 3, TGFR-3, Betaglycan, Transforming growth factor beta receptor III, TGF-beta receptor type III, TGFBR3

## **Dilution**

WB~~1/500 - 1/2000 E~~1/10000

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

TGFBR3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **TGFBR3 Antibody - Protein Information**



## Name TGFBR3 (HGNC:11774)

#### **Function**

Cell surface receptor that regulates diverse cellular processes including cell proliferation, differentiation, migration, and apoptosis (PubMed:<a

href="http://www.uniprot.org/citations/12958365" target="\_blank">12958365</a>, PubMed:<a href="http://www.uniprot.org/citations/19416857" target="\_blank">19416857</a>). Initiates BMP, inhibin, and TGF-beta signaling pathways by interacting with different ligands including TGFB1, BMP2, BMP5, BMP7 or GDF5 (PubMed:<a

href="http://www.uniprot.org/citations/18184661" target="\_blank">18184661</a>). Alternatively, acts as a cell surface coreceptor for BMP ligands, serving to enhance ligand binding by differentially regulating BMPR1A/ALK3 and BMPR1B/ALK6 receptor trafficking (PubMed:<a href="http://www.uniprot.org/citations/19726563" target="\_blank">19726563</a>). Promotes epithelial cell adhesion, focal adhesion formation and integrin signaling during epithelial cell spreading on fibronectin (PubMed:<a href="http://www.uniprot.org/citations/22562249" target="\_blank">22562249</a>). By interacting with the scaffolding protein beta-arrestin2/ARRB2, regulates migration or actin cytoskeleton and promotes the activation of CDC42 as well as the inhibition of NF-kappa-B (PubMed:<a

href="http://www.uniprot.org/citations/19416857" target="\_blank">19416857</a>, PubMed:<a href="http://www.uniprot.org/citations/19325136" target="\_blank">19325136</a>). In gonadotrope cells, acts as an inhibin A coreceptor and regulates follicle-stimulating hormone (FSH) levels and female fertility (By similarity). Plays a role in the inhibition of directed and random cell migration in epithelial cells by altering the actin cytoskeletal organization (PubMed:<a href="http://www.uniprot.org/citations/19416857" target="\_blank">19416857</a>). Participates in epithelial-mesenchymal transformation (EMT) upon binding to BMP2 or TGFB2, by activating the PAR6/SMURF1/RHOA pathway (By similarity).

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein. Secreted {ECO:0000250|UniProtKB:P26342}. Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P26342}. Note=Exists both as a membrane-bound form and as soluble form in serum and in the extracellular matrix. {ECO:0000250|UniProtKB:P26342}

## **TGFBR3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture






Figure 1: Western blot analysis using TGFBR3 mAb against human TGFBR3 recombinant protein. (Expected MW is 44.1 kDa)

17-

11-




Figure 2: Western blot analysis using TGFBR3 mouse mAb against Jurkat (1), HeLa (2), MCF-7 (3), F9 (4), SK-N-SH (5), and NIH3T3 (6) cell lysate.

# **TGFBR3 Antibody - References**

1.Mol Biol Cell. 2011 May;22(9):1463-72. 2.Zhongguo Fei Ai Za Zhi. 2010 May;13(5):451-7.



