

CD154
Purified Mouse Monoclonal Antibody
Catalog # AO2711a**Specification**

CD154 - Product Information

Application	WB, IHC, ICC, E
Primary Accession	P29965
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2a
Calculated MW	29.3kDa KDa

Immunogen

Purified recombinant fragment of human CD154 (AA: extra 47-261) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

CD154 - Additional Information

Gene ID 959

Other Names

CD40LG; IGM; IMD3; TRAP; gp39; CD40L; HIGM1; T-BAM; TNFSF5; hCD40L

Dilution

WB~~ 1/500 - 1/2000

IHC~~1:100~500

ICC~~N/A

E~~ 1/10000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD154 is for research use only and not for use in diagnostic or therapeutic procedures.

CD154 - Protein Information

Name CD40LG

Synonyms CD40L, TNFSF5, TRAP

Function

Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:<a

[1280226](http://www.uniprot.org/citations/1280226), PubMed: [31331973](http://www.uniprot.org/citations/31331973)). Costimulates T-cell proliferation and cytokine production (PubMed: [8617933](http://www.uniprot.org/citations/8617933)). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed: [8617933](http://www.uniprot.org/citations/8617933)). Induces the activation of NF-kappa-B (PubMed: [15067037](http://www.uniprot.org/citations/15067037), PubMed: [31331973](http://www.uniprot.org/citations/31331973)). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed: [15067037](http://www.uniprot.org/citations/15067037)). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed: [15067037](http://www.uniprot.org/citations/15067037)). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell surface

Tissue Location

Specifically expressed on activated CD4+ T- lymphocytes

CD154 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD154 - Images




Figure 1: Black line: Control Antigen (100 ng); Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng)




Figure 2: Western blot analysis using CD154 mAb against human CD154 (AA: extra 47-261) recombinant protein. (Expected MW is 50 kDa)




Figure 3: Western blot analysis using CD154 mAb against HEK293 (1) and CD154 (AA: extra 47-261)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 4:Flow cytometric analysis of Ramos cells using CD154 mouse mAb (green) and negative control (red).

CD154 - References

1.Hematology. 2016 Apr;21(3):187-92. 2.Mol Cell Probes. 2015 Dec;29(6):335-342.