

C2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10662a**Specification**

C2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P06681
Other Accession	NP_000054
Reactivity	Human, Hamster
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	83268
Antigen Region	147-176

C2 Antibody (N-term) - Additional Information**Gene ID** 717**Other Names**

Complement C2, C3/C5 convertase, Complement C2b fragment, Complement C2a fragment, C2

Target/Specificity

This C2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 147-176 amino acids from the N-terminal region of human C2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

C2 Antibody (N-term) - Protein Information**Name** C2**Function** Component C2 which is part of the classical pathway of the complement system is

cleaved by activated factor C1 into two fragments: C2b and C2a. C2a, a serine protease, then combines with complement factor C4b to generate the C3 or C5 convertase.

Cellular Location


Secreted.

C2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C2 Antibody (N-term) - Images



C2 Antibody (N-term) (Cat. #AP10662a) western blot analysis in CHO cell line lysates (35ug/lane). This demonstrates the C2 antibody detected the C2 protein (arrow).

C2 Antibody (N-term) - Background

Component C2 is a serum glycoprotein that functions as part of the classical pathway of the complement system. Activated C1 cleaves C2 into C2a and C2b. The serine proteinase C2a then combines with complement factor 4b to create the C3 or C5 convertase. Deficiency of C2 has been reported to associated with certain autoimmune diseases and SNPs in this gene have been associated with altered susceptibility to age-related macular degeneration. This gene localizes within the class III region of the MHC on the short arm of chromosome 6. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described in publications but their full-length sequence has not been determined.

C2 Antibody (N-term) - References

Hu, M., et al. Pharmacogenet. Genomics 20(10):634-637(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Liu, X., et al. Retina (Philadelphia, Pa.) 30(8):1177-1184(2010)
Han, S., et al. Hum. Immunol. 71(7):727-730(2010)
Ishii, Y., et al. J. Immunol. 151(1):170-174(1993)