

C17orf59 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11018c**Specification**

C17orf59 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q96GS4
Other Accession	Q66H43 , Q9D6W8 , Q3SX20 , NP_060092.2
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	186-215

C17orf59 Antibody (Center) - Additional Information**Gene ID** 54785**Other Names**

Uncharacterized protein C17orf59, C17orf59

Target/Specificity

This C17orf59 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 186-215 amino acids of human C17orf59.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C17orf59 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

C17orf59 Antibody (Center) - Protein Information**Name** BORCS6 ([HGNC:25939](#))**Function** As part of the BORC complex may play a role in lysosomes movement and localization

at the cell periphery. Associated with the cytosolic face of lysosomes, the BORC complex may recruit ARL8B and couple lysosomes to microtubule plus-end-directed kinesin motor.

Cellular Location


Lysosome membrane.

C17orf59 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C17orf59 Antibody (Center) - Images



C17orf59 Antibody (Center) (Cat. #AP11018c) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the C17orf59 antibody detected the C17orf59 protein (arrow).

C17orf59 Antibody (Center) - References

Kimura, K., et al. Genome Res. 16(1):55-65(2006)
Suzuki, Y., et al. Gene 200 (1-2), 149-156 (1997) :